

**NAMIBIA NATIONAL CANCER REGISTRY (NNCR)
CANCER INCIDENCES IN NAMIBIA 2010 - 2014**

Data collected and entered by the Namibia National Cancer Registry (NNCR)

Dr Annelle Zietsman

Sr Johanna Pontac

Sr Sophy Lawrence

Ms Lizette Horn

Mrs Lizelle van Schalkwyk

As National Director: Namibia National Cancer Registry

Mrs Reinette Koegelenberg (previous director)

Mr Rolf-Jürgen Hansen (current director)

Data analysis and report written by

Mr Henri Carrara

Consultant Analytical Epidemiologist (RSA)

Report reviewed by

Dr Maxwell Parken

African Cancer Registry Network (AFCRN)

Supported by



Published February 2017

FOREWORD

NAMIBIA NATIONAL CANCER REGISTRY REPORT 2010 - 2014

The Namibia National Cancer Registry is a combined project between the Ministry of Health and Social Services (MOHSS) and the Cancer Association of Namibia (CAN) with data analysis done by IARC.

This collaboration started in 1995 and this report is the fifth published report.

I would like to congratulate and thank everyone involved in this project - the doctors and nurses from MOHSS collecting data; the CAN staff doing data entry and follow-up; IARC doing data- analysis and the sponsors involved in producing the report. Without the hard work of each individual this very important research project would not be possible.

This information is essential for the action that must be taken by the Namibian National Cancer Control Program – another collaborative program to start soon.

In our collective battle against cancer our vision and mission is to:

1. Prevent cancers - create awareness, provide education about healthy lifestyle, promote HPV vaccination
2. Promote early detection - screening
3. Provide information about the most common cancers in Namibia
4. Provide adequate medical treatment for all Namibian cancer patients
5. Provide palliative and supportive care for all Namibians

This battle can only be fought by a multidisciplinary team which include MOHSS, CAN, all healthcare professionals, the government of Namibia, the media, churches, NGOs and every Namibian citizen.

With sincere gratitude

Annelle Zietsman

Dr. A.B. May Cancer Care Centre – Windhoek Central Hospital

MOHSS

A SPECIAL THANK YOU

A special word of thanks to Morten Ervik of IARC for help answering many questions relating to CanReg5 software.

The Cancer Association of Namibia would like to thank all medical practitioners, medical and treatment establishments and volunteers who have supported data collection.

We would also like to thank the African Cancer Registry Network (AFCRN) for their continued support on cancer reporting in not only Namibia, but Africa as a whole.

CONTENTS

Message from the Cancer Association of Namibia (CAN) and Namibia National Cancer Registry (NNCR)P6

SUMMARY.....P7

INTRODUCTION.....P8

Background of the Namibia National Cancer Registry

History of the registry

Location of registry

Source of finance

Previous publications

Aims of the Namibia National Cancer Registry

Namibia: A brief introduction

MATERIALS AND METHODSP12

Data collection by the Namibia National Cancer Registry

Sources of information for the registry

Death registry

Methods of registration

Data management

Data analysed in this report

Years analysed

Variables examined

Population denominators

Data analysis and presentation

RESULTSP14

Overall burden of cancer in Namibia 2010-2014

Leading cancers among males

Prostate cancer

Kaposi sarcoma

Non-Hodgkin lymphoma

Eye cancer

Lung/trachea/bronchus cancer

Other notable cancers in males

Leading cancers among females

Breast cancer

Cervix cancer

Kaposi sarcoma

Eye cancer

Ovarian cancer

Other notable cancers in females

Childhood cancers

Basis of diagnosis of cancers

DISCUSSIONP23

Limitations

Increasing burden of cancer in Namibia

Cancers associated with Human Immunodeficiency Virus infection

Prostate cancer

Tobacco and alcohol-related cancers

Female-specific cancers

Colon cancer

Basal and squamous cell skin cancers

Cancers among children

RECOMMENDATIONSP26

Accurate collection of regional data

Provision of population statistics by ethnic group

CONCLUSIONP26

FIGURES AND TABLES

Figure 1: Map of Namibia: regions and population	P8
Figure 2: Map of Namibia: regional in SADC situation.....	P8
Figure 3: NAMIBIA.....	P9
Figure 4: NAMIBIA.....	P9
Figure 5: CAN	P10
Figure 6: A.B. MAY	P10
Figure 7: NOC	P11
Figure 8: Population of Namibia (2012, middle of analysis period).....	P14
Figure 9: Bar Chart showing Top 10 Cancers in Namibia (2010-2014), excluding C44, Male	P15
Figure 10: Age-specific Incidence rates per 100 000 in 2010-2014, Prostate (C51)	P16
Figure 11: Age-specific Incidence rates per 100 000 in 2010-2014, Kaposi sarcoma (C46).....	P16
Figure 12: Age-specific Incidence rates per 100 000 in 2010-2014, Non-Hodgkin lymphoma (C82-85, C96) ...	P17
Figure 13: Age-specific Incidence rates per 100 000 in 2010-2014, Eye (C69)	P18
Figure 14: Age-specific Incidence rates per 100 000 in 2010-2014, Trachea, bronchus and lung (C33-34)	P18
Figure 15: Bar Chart showing Top 10 Cancers in Namibia (2010-2014), excluding C44, Female.....	P15
Figure 16: Age-specific Incidence rates per 100 000 in 2010-2014, Breast (C50).....	P20
Figure 17: Age-specific Incidence rates per 100 000 in 2010-2014, Cervix uteri (C53)	P20
Figure 18: Age-specific Incidence rates per 100 000 in 2010-2014, Kaposi sarcoma (C46).....	P21
Figure 19: Age-specific Incidence rates per 100 000 in 2010-2014, Eye (C69)	P22
Figure 20: Age-specific Incidence rates per 100 000 in 2010-2014, Ovary (C56)	P22
Table 1: NAMBIA (2010-2014) Annual incidence per 100 000 by age group - Male	P29
Table 2: NAMBIA (2010-2014) Annual incidence per 100 000 by age group - Female.....	P29
Table 3: NAMBIA (2010-2014) Cancer frequencies by ethnic group – Males.....	P30
Table 4: NAMBIA (2010-2014) Cancer frequencies by ethnic group – Females	P30
Table 5: NAMBIA (2010-2014) Cancer frequencies by region – Males.....	P31
Table 6: NAMBIA (2010-2014) Cancer frequencies by region – Females	P31
Table 7: NAMBIA (2010-2014) Basis of diagnosis of cancers – Males	P32
Table 8: NAMBIA (2010-2014) Basis of diagnosis of cancers - Females.....	P32
Appendix: NAMBIA Annual incidence per 100 000 by age group and sex for single years	P33

Appendix: C15 editorial tables P38

Appendix: Data quality indicators Namibia National Cancer Registry P42

MESSAGE FROM THE CANCER ASSOCIATION OF NAMIBIA (CAN) AND NAMIBIA NATIONAL CANCER REGISTRY (NNCR)

Founded in 1968 by the late Dr A Bernard May, the Cancer Association of Namibia (WO30), an incorporated association not for gain registered with the Ministry of Health and Social Services and Ministry of Trade and Industry, has continuously strived to better the conditions related to the fight against, and support of those afflicted by, cancer in our country.

The Association remains actively engaged in creating awareness, educating and supporting Namibians in the fight against cancer. A key element of the war on cancer is supporting data collection and presentation of a cancer incidence report in Namibia.

Commitment on all levels and from all stakeholders remains pivotal in making an impact on this journey of change.

Change plays a cardinal role, because a change in attitude, breaking the stigma and developing better national plans to combat diseases, is part of our mission to rid our communities of non-communicable diseases like cancer.

How do we eradicate it, if we do not know what we are up against?

The Cancer Association of Namibia, as administrator of the Namibia National Cancer Registry, is therefore proud to present the fifth 'Cancer Incidence in Namibia Report (2010 – 2014)'. This document should ideally be read along with the previous published reports (1995 – 1998; 2000 – 2005; and 2006 – 2009). From commentary, you will quickly realize that there is still much work to be done to ensure we provide data of a world-class standard from Namibia in terms of cancer reporting. It is also evident that as time goes by, case reporting, active case finding and case updating does enjoy more attention.

Data collection is a very time-consuming and a costly process and, as a welfare organisation, we underscore the urgency of public and stakeholder engagement – not only in data support, but for funding as well. Since the Association remains committed to fighting cancer and to making an impact on our communities, the national cancer registry must remain a priority at all costs!

I would like to thank my predecessors for establishing a strong foundation to develop the NNCR further, while the staff of the NNCR – in particular, Ms Lizette Horn – deserves applause for her dedication and commitment to the cause. We acknowledge and thank the staff of the Dr AB May Cancer Care Centre – Dr Anelle Zietsman and her team; as well as all other data collection stakeholders nationwide. A special word of thanks to Prof Max Parkin of the African Cancer Registry Network (AFCRN) for their commitment to data collection not only in Namibia, but Africa as a whole.

In conclusion, I trust that this publication will serve as a guide to all parties involved in decision-making, advocacy and responsible citizenry (both private and corporate) to support the advocacy on cancer and in so doing, reduce the incidence of this disease that remains the leading cause of death worldwide.

Rolf-Jürgen Hansen
Chief Executive Officer
Cancer Association of Namibia

SUMMARY

This report summarises the results of an analysis of cancers reported to the Namibia National Cancer Registry from 2010 to 2014.

A total of 11 248 malignant neoplasms were recorded among the Namibian population during this five-year period, of which 5 125 were in males (45.6%) and 6 123 in females (54.4%). With 'other skin' neoplasms excluded, there were 10 559 people with malignant neoplasms; 4 787 males (45.3%) and 5 772 females (54.7%). Of this number, 353 (3.3%) were in children below the age of 15.

Overall, the incidence of all cancers including non-melanoma skin in males was 153.8 per 100 000 and 146.3 per 100 000 in females. Excluding non-melanoma skin cancers, the incidence of cancer overall in males was 144.2 per 100 000 and was 138.7 per 100 000 in females. This represents an increase in incidence in both sexes compared to the previous reporting period (2005–2009) while the ranking of the most common types of cancer remained similar with the exception of a notable increase in the HIV-associated cancers (Kaposi sarcoma, Non-Hodgkin lymphoma and Eye cancer).

The most common cancers in Namibia by gender: 2010-2014

Cancer in males	Proportion of all cancers (%)	ASR / 100 000	Cancer in females	Proportion of all cancers (%)	ASR / 100 000
Prostate	23.6	37.6	Breast	27.4	40.2
Kaposi sarcoma	18.4	22.0	Cervix	19.4	27.2
Non-Hodgkin lymphoma	4.8	6.3	Kaposi sarcoma	8.8	10.1
Eye	5.0	6.2	Eye	4.0	4.8
Lung/trachea/bronchus	3.4	5.9	Ovary	3.0	4.3
Larynx	3.1	5.3	Non-Hodgkin lymphoma	2.8	3.4
Colon	2.7	4.2	Corpus uteri	2.1	3.3
Liver	2.6	4.0	Lung/trachea/bronchus	1.8	2.9

This report describes the overall occurrence of cancer and the leading cancers in Namibia. This data continues to form a basis for future research into risk factors for cancers in the Namibian population and for program planning and advocacy. It is recommended that cancer be made a reportable condition in Namibia to improve these cancer statistics.

INTRODUCTION

The World Health Organisation (WHO) estimates that more than 14.1 million new cases of cancer occur annually and that more than half of these occur in developing countries (GLOBOCAN 2012, IARC). Cancer is the most common cause of death in developed countries. It is difficult to precisely estimate the burden of cancer mortality in developing countries due to lack of and/or the poor quality of available data. However, it is estimated that 10% of deaths in developing countries are due to cancer and given the aging of these populations, cancer incidence is increasing (WHO, 2002).

A cancer registry provides a framework for assessing and measuring the impact of cancer in communities and is important in the allocation of health resources. It is an essential surveillance device for measuring the effectiveness of any cancer control program. The data produced furthermore provides a starting point for research into cancer causes and prevention.

BACKGROUND OF THE NAMIBIA NATIONAL CANCER REGISTRY

History of the registry

The Namibia National Cancer Registry (NNCR) was started in 1995, when Rössing Uranium mine, in co-operation with the Namibian Ministry of Health (Oncology clinic) and the Cancer Association of Namibia, collected all cancer cases reported to the Windhoek state pathology laboratory and the single existing private pathology laboratory from 1979 to 1994. From 1995 onwards, the Namibia National Cancer Registry commenced active registration of both pathology-based and clinical cases for four sentinel regions (Erongo, Oranjemund, Oshakati and Khomas). Cases that are diagnosed in South Africa are re-routed to the Namibian registry via a network of registries, which are technically supported by the International Agency for Research on Cancer (IARC). The Namibia National Cancer Registry aims to provide a database of information that will ultimately lead to improved cancer prevention and control among the Namibian population.

The Namibia National Cancer Registry is a population-based registry.

Location of registry

The registry is based at the premises of the Cancer Association of Namibia (90 John Meinert Street, Windhoek-West, WINDHOEK). There is a separate office with one computer linked to a server; this is used for data entry. Transport, technical and administration support are provided by the Association, as required.

The registry is co-managed by Mr R. Hansen, CEO of the Cancer Association of Namibia. The Technical Leader is Ms L. Horn (Cancer Association of Namibia) and Dr. A. Zietsman and the staff of the Dr A.B. May Cancer Care Centre guide with data input and patient identification.

The registry covers the entire population of Namibia enumerated as 2,104,900 persons at the 2011 census. The map of Namibia (Figure 1) shows the 13 regions and the population of each.




Figure 1: Map of Namibia: regions and population




Figure 2: Map of Namibia: regional in SADC situation



Figure 3: NAMIBIA: Namib Desert



Figure 4: NAMIBIA: Key Towns



Figure 5: CANCER ASSOCIATION OF NAMIBIA



Figure 6: DR A.B. MAY CANCER CARE CENTRE - State cancer care



Figure 7: NAMIBIA ONCOLOGY CENTRE – Private cancer care

Source of finance

The International Association of Cancer Registries (IARC) and African Cancer Registry Network (AFCRN) cover some costs pertaining to the registry. The Cancer Association of Namibia (WO30) funds all other registry activities.

Previous publications

There are four previously published reports in this series (1989-1994, 1995-1998, 2000-2005; and 2006-2009).

- Koehler S. Namibia's cancer atlas: 1989-1994. Namibia National Cancer Registry 1997.
- Stein L, Sitas F, Zietsman A, Gariseb B, Rautenbach N, Pontac J and Johannesson I. Cancer in Namibia 1995 - 1998, Namibia Cancer Registry, Windhoek 1999.
- Carrara H, Stein L, Sitas F, Zietsman A, Gariseb B, Rautenbach N, Pontac J, Johannesson I and Ferlay J. Cancer in Namibia 2000-2005, Namibia Cancer Registry, Windhoek 2009.
- Carrara H, Stein L, Sitas F, Zietsman A, Gariseb B, Rautenbach N, Pontac J, Rooi T, and Taapopi L. Cancer in Namibia 2006-2009, Namibia Cancer Registry, Windhoek 2011.

A number of other publications and conference abstracts have been prepared.

- The NNCR has already contributed to the body of literature on Cancer in Africa (Parkin et al., 2003; Parkin et al., 2008; Sitas et al., 2008, Ferlay et al, 2010 – the WHO/IARC database GLOBOCAN 2012).
- ASB van Dyk, RN, LF Small, A Zietsman, The Pain Experience and its management in cancer patients during hospitalization (in Namibia) Health SA Gesondheid Vol.5 No.4 – 2000.

Aims of the Namibia National Cancer Registry

The NNCR aims to provide a database of information that will ultimately lead to improved cancer prevention and control among the Namibian population. The main goals of the NNCR are:

- To monitor the incidence of cancer among the Namibian population on an ongoing basis;
- To use observed cancer trends to predict future cancer patterns in Namibia;
- To provide information on the burden of cancer in different regions and among different ethnic groups in Namibia;
- To monitor the effects of cancer prevention programs, early detection or screening, treatment and palliative care;

- To provide information that will serve as a basis for research into cancer causes specific to the Namibian population.

Namibia: A brief introduction

Namibia, a southern African country shares borders with Angola and Zambia to the north, Botswana to the east, and South Africa to the south. It gained independence from South Africa in 1990 and its capital city is Windhoek. With only 2.5 inhabitants per square kilometre it is one of the most sparsely populated countries in the world.

Namibia is one of Africa's most developed and stable countries. Tourism and mining form the backbone of Namibia's economy.

The majority of the Namibian population is Black African — mostly of Ovambo ethnicity, which forms about half of the population — and is concentrated in the north of the country. There are also the Herero and Himba people who speak a similar language. In addition to the Black African majority, there are large groups of Khoisan (the Nama and San). Other Black Africans are descendants of refugees from Angola. There are also two smaller groups of people with mixed racial origins, called "Coloureds" and "Basters", who together make up 6% of the population. Whites of Portuguese, Dutch, German, British and French ancestry also make up about 6% of the population. Most of Namibian Whites and nearly all those of mixed race are Afrikaans speakers and share similar origins, culture, religion and genealogy as the White and Coloured populations of neighbouring South Africa. A smaller proportion of Whites, approximately 30 000, trace their origins directly back to German settlers and maintain German cultural and educational institutions. Figure 1 shows the map of Namibia and the geographic regions used in this report.

MATERIALS AND METHODS

Data collection by the Namibia National Cancer Registry

The NNCR attempts to record for each cancer patient, first name and surname, date of birth or approximate age, I.D. number, gender, marital status, ethnic origin, current residential address, place of birth, tobacco and alcohol use, reproductive history (for females), date of cancer diagnosis, histology number, tumour site and histology, clinical stage of disease, basis of diagnosis, treatment given, vital status, and date and cause of death.

Until late 2010 data from the cancer registration form were entered into CanReg4 Software (Cooke et al., 2001; Cook et al., 2005). The data were migrated to CanReg5 (<http://canreg.iarc.fr>) in late 2010 and the data were validated and duplicates were identified using check and conversion programs for cancer registries (Ferlay et al., 2005). Tumour site and histology were coded using ICD-0-3 (2) (Fitz et al., 2001) at the time of data entry, but were converted to ICD-10 for the tabulation of cases. Since 2010 ICD-10 has been used to code cancer sites.

Sources of information for the registry

There are two large state hospitals in Windhoek: Central Hospital (which includes the Dr A.B. May Cancer Care Centre) and Katutura Hospital, and a referral hospital in each region of the country. A cancer centre recently opened at Oshakati Intermediate Hospital. In addition, there are five private hospitals in Windhoek (Roman Catholic, Rhino Park, Medi-Clinic, Paramount and Lady Pohamba).

During 2015 the Namibia Oncology Centre (NOC) – a specialized private cancer treatment facility – opened its doors; the first of its kind for Namibia.

At present, primary data collection takes place at the Dr A.B. May Cancer Care Centre and the Namibia Oncology Centre to which all cancer patients are supposed to be sent for assessment and possible treatment if private or state. Paramount Hospital and GVI Oncology in Namibia also serve as data collection and patient referral points.

There are two pathology laboratories in the country: the state run National Institute of Pathology (NIP) and the private run, PathCare. Some information on cases from Namibia is also received from a laboratory in Cape Town (Dr Holm).

The pathology reports from NIP include demographic data, but place of residence is noted in less than one third of these. The print-out received from PathCare does not include personal details, only the age and sex of the subject. For this reason, completeness of data remains a great concern, and the registry and Cancer Association of Namibia have been actively engaged with the Ministry of Health and Social Services to have cancer declared a reportable disease. In addition, active case finding throughout the vast country should be urgently implemented.

There is no service for clinical haematology.

Death registry

Civil registration of deaths, by cause, is the responsibility of the Ministry of Home Affairs. Permission to access to copies of certificates for cancer cases was granted from 2016 only. This is because a new electronic registry and databank is finally in place. The quality of information as to cause of death is not known.

Methods of registration

Case finding relies upon receipt of registration forms, completed by nursing staff in the admissions unit of the Dr A.B. May Cancer Centre, NOC and copies of pathology reports, as described above. Currently, there is no active case finding and minimal follow-up in Namibia, unless conducted by staff of the Cancer Association of Namibia on an ad-hoc basis. Lack of funding in this regard hampers this part of the project, as only the Cancer Association of Namibia and some donors who agree on the urgency of data pertaining to incidence reports cover costs relating to the NNCR.

Data management

The registry is housed at the Cancer Association of Namibia in a dedicated office. A desktop computer is used for data entry employing the specialised cancer registration software running in a Microsoft Windows environment. CanReg5 is employed.

Prior to data analysis, the data were exported as a comma-separated-values dataset and these data were checked using the IARC check and conversion programs. This generated a set of data that were manually checked by registry staff who updated the data in CanReg5 for analysis.

Data analysed in this report

Years analysed

The data collected by the NNCR consists of 31 788 cancer cases recorded from 1969 to 2014. Between 1969 and 1994, the number of cases recorded per year ranged from 1 to 402. Data from 1989 to 1994 was analysed in the first report (Koehler, 1997). Active registration of both pathology-based and clinical cases was only started in 1995. A total of 5 141 cases¹ were recorded in the period from 1995 to 1998, approximately 1 285 cases per year. In 1999 only 534 cases were recorded, thus these data were not included in any of the reports. The subsequent report covered the period from 2000 to 2005 and included 5 009 cases, an average of approximately 834 cases per year (Carrara et al, 2009). This was followed by a report covering the period 2006 to 2009 during which a total of 6 363 cases, an average of 1 591 cases each year, was analysed (Carrara et al, 2011). The current report represents the most complete and reliable reflection of cancer incidence in the Namibian population between 2010 and 2014. A total of 11 248 malignant neoplasms were recorded among the Namibian population during this five-year period, an average of 2 250 cancers per year, an increase of almost 40% on the previous reporting period.

Variables examined

Variables used in the analysis of the data from 2010 to 2014 included age (calculated from date of birth), gender, current region of residence, ethnicity, date of cancer diagnosis, tumour site and histology, and basis of diagnosis. The variables describing tobacco and alcohol use, as well as female reproductive history, have improved since the previous report and will be examined further in another publication.

Population denominators

The population at risk for the period 2010-2014 used for incidence calculations was the mid-period data for 2012. This was provided by the Namibian Central Bureau of Statistics based on projected population estimates modelled on the census data of 2011 (Republic of Namibia, Central Bureau of Statistics, 2014). The annual population at risk for the whole of Namibia for the period 2012 is shown in Figure 8.

¹ Previous reports in this series may contain minor differences in numbers resulting from ongoing correction of the dataset.




Figure 8: Population of Namibia (2012, middle of analysis period)

Data analysis and presentation

The data were analysed in Cape Town, South Africa, employing CanReg5 software running on Mackintosh IOS platform. Incidence rates were calculated for the whole of the Namibian population in 5-year age groups. The upper age group was set at 80 and above. Age standardisation was performed by the direct method, using the world standard population (Parkin et al., 1997).

Cancer incidence by sex and 5-year age group (Tables 1 and 2) were generated using CanReg5, IARC. These data are disaggregated by year and shown in the appendix. Frequencies of cancer by ethnic group (Tables 3 and 4) and region (Tables 5 and 6) were generated manually in CanReg5 and collated into tabular format using Microsoft Excel.

Occurrence rates could only be calculated for males and females for the whole of Namibia, as population denominators in 5-year age groups were not available for different ethnic groups, or geographic regions by sex. Consequently only frequency tables are presented to summarise the available results for ethnic group and geographic region. Rate calculations for 'all cancers' and ranking of cancers exclude 'other skin' (ICD10-C44), which is standard practice in cancer registry reports, and allows for international comparisons.

The appendix also contains a copy of the Cancer Incidence in Five Continents editorial tables for major diagnosis groups for the readers' further information and to allow international comparisons.

The appendix further contains a table of data quality indicators, also utilising the major diagnosis groupings by sex, here %MV represents the proportion of cases that are microscopically verified.

RESULTS

A total of 11 248 malignant neoplasms were recorded among the Namibian population during this five-year period under review, of which 5 125 occurred in males (45.6%) and 6 123 occurred in females (54.4%). The corresponding average number of cases reported annually was 2 250 overall, 1 025 in males and 1 226 in females. With 'other skin' excluded, the data set consisted of 10559 malignant neoplasms, 4 787 occurred in males (45.3%) and 5 772 occurred in females (54.7%). Of this number, 353 (3.3%) recorded cases were in children below the age of 15. Thus an average of 71 cases was reported annually in children. Ninety-five percent of all cases were verified histologically.

Tables 1 and 2 show the frequencies, incidence rate per 100 000 in 5-year age groups, crude rate, proportions of all cancers, cumulative rates (0-64), cumulative rates (0-74) and age standardised rates (ASR) by site and gender for the period 2010 to 2014. Tables 3 and 4 show cancer frequencies and proportions by ethnic group for males and females respectively. Tables 5 and 6 show cancer frequencies and proportions by geographic region of residence for males and females, respectively. Tables 7 and 8 show the number and proportions for the basis of cancer diagnosis by sex. Appendix 1 shows the cancer frequencies by site and gender in single calendar years of observation for 2010, 2011, 2012, 2013 and 2014. This shows less variation in this 5-year reporting period compared to the previous 4-year reporting period (2005-2009), but the incidence of cancer was still substantially higher in 2014 than in each of the years from 2010-2013. Of all the data collected thus far, the numbers reported here are likely to be the most accurate reflection of the annual burden of cancer in Namibia.

Overall burden of cancer in Namibia 2010-2014

A total of 11 248 malignant neoplasms were recorded among the Namibian population during this five-year period, of which 5 125 were in males (45.6%), and 6 123 in females (54.4%). With ‘other skin’ neoplasms excluded, there were 10 559 people with malignant neoplasms; 4 787 males (45.3%) and 5 772 females (54.7%). Of this number, 353 (3.3%) were in children below the age of 15.

Overall, the incidence of all cancers including non-melanoma skin in males was 153.8 per 100 000 and in females it was 146.3 per 100 000. Excluding non-melanoma skin cancers, the incidence of cancer overall in males was 144.2 per 100 000 and in females it was 138.7 per 100 000. This represents an increase in incidence in both sexes compared to the previous reporting period (2005-2009) when the ASR in males and females excluding non-melanoma skin was 90.2 per 100 000 and 100.7 per 100 000 respectively.²

The ranking of the most common types of cancer remained similar with the exception of a notable increase in the HIV-associated cancers (Kaposi sarcoma, Non-Hodgkin lymphoma and Eye cancer).

Leading cancers among males

The most frequent cancer among men between 2010 and 2014 was prostate cancer (23.6%, ASR 37.6), followed by Kaposi sarcoma (18.4%, ASR 22.0), Non-Hodgkin lymphoma (4.8%, ASR 6.3), cancers of the eye (5.0%, ASR 6.2), lung / trachea / bronchus (3.4%, ASR 5.9), larynx (3.1%, ASR 5.3), colon (2.7%, ASR 4.2), liver (2.6%, ASR 4.0), mouth (2.4%, ASR 3.9) and oesophagus (2.3%, ASR 3.9).²




Figure 9: Bar Chart showing Top 10 Cancers in Namibia (2010-2014), excluding C44, Male

² The ranking of top cancers in the report is based on age standardized rates, while the ranking depicted in the bar chart is on absolute numbers.

Prostate cancer

Cancer of the prostate was the most common cancer among males during the period beginning in 2010 and ending in 2014. A total of 1 128 cases were reported, comprising 23.6% of all male cancers. The annual ASR for prostate cancer was 37.6 per 100 000, an almost two-fold increase compared to the previous reporting period (ASR 23.5). Incidence increased with age from 1.6 per 100 000 at age 30-34 to 546.5 per 100 000 for men aged 75-79.




Figure 10: Age-specific Incidence rates per 100 000 in 2010-2014, Prostate (C51)

Cancer of the prostate was the leading cancer for the Tswana (35.0%), Baster (33.9%), Herero (29.1%), Coloured (22.1%), White (19.9%), Nama (15.5%) and San/Bushman (11.8%) ethnic groups.

This cancer was among the five leading male cancers in almost all regions and was the most frequent cancer in males in the Oshana (32.2%), Khomas (31.4%), Kunene (25.9%), Omaheke (25.0%), Hardap (19.0%), Ohangwena (18.1%) and Kharas (18.0%) regions.

Kaposi sarcoma

A total of 882 cases of Kaposi sarcoma (KS) were reported from 2010 to 2014, comprising 18.4% of all cancer cases in males during this five-year period. The overall ASR for KS was 22.0 per 100 000, having increased from 16.3 per 100 000 in the previous reporting period. The highest incidence occurred among males aged between 40 and 44 years (ASR 62.4 per 100 000).




Figure 11: Age-specific Incidence rates per 100 000 in 2010-2014, Kaposi sarcoma (C46)

On average, during this five-year period, KS was the leading cancer among males in the Ovambo (24.1%), Caprivi (19.8%) and Kavango (10.1%) ethnic groups. KS was furthermore among the five leading cancers in males in all ethnic groups examined, except for the White, Baster, Coloured, San / Bushman and Nama ethnic groups. It should be noted that very few cancers were recorded in the San / Bushman and Tswana ethnic groups overall.

KS was the leading cancer in males in almost all regions of Namibia; Kavango (34.3%), Caprivi (31.6%), Oshikoto (26.1%), Omusati (23.9%), Otjozondjupa (23.7%), Ohangwena (18.1%), Erongo (14.2%) and Hardap (9.8%) regions, and among the five leading cancers in other regions.

Non-Hodgkin lymphoma

Non-Hodgkin lymphoma (NHL) ranked third among cancers in Namibian males for the period 2010 to 2014 (231 cases reported, comprising 4.8% of all cancers). The ASR for NHL in males was 6.3 per 100 000, having almost doubled since the previous reporting period. Within age groups, the annual incidence per 100 000 tended to increase with age from 1.5 in males aged 0-4, up to 23.6 in males aged 70-74 years.




Figure 12: Age-specific Incidence rates per 100 000 in 2010-2014, Non-Hodgkin lymphoma (C82-85, C96)

NHL was among the top five cancers in males of the Caprivi (7.4%), White (6.0), San (5.9%), Herero (5.5%), Kavango (4.8%) and Ovambo (4.8%) ethnic groups.

NHL was the third most common cancer among males living in the Oshikoto (9.0%), Otjozondjupa (8.1%), Omusuti (6.1%) and Caprivi (3.9%) regions and among the top ten cancers in all the other regions except the Ohangwena Region.

Eye cancer

Cancer of the eye was the fourth most common cancer in this reporting period, 2010 to 2014. A total of 241 cases were reported, comprising 5.0% of all male cancers. The ASR for cancer of the eye was 6.3 per 100 000; up from 2.3 in the previous reporting period. The incidence increased with age, from 2.4 per 100 000 in those younger than 5 years, up to 19.3 per 100 000 in those aged 65-69 years.




Figure 13: Age-specific Incidence rates per 100 000 in 2010-2014, Eye (C69)

Cancer of the eye was among the five most common cancers among the Damara (7.3%), Ovambo (6.1%), Herero (5.8%) and Baster (5.3%) ethnic groups.

Cancer of the eye was the third most frequent cancer in males living Oshana (8.2%), Khomas (6.7%) and the Kavango (4.3%) regions, and the fifth most common cancer in the Omasuti (4.5%) Region.

Lung/trachea/bronchus cancer

Cancer of the lung/trachea/bronchus was the fifth most common cancer in Namibia in this reporting period. A total of 161 cases were reported, comprising 3.4% of all male cancers. The ASR for cancer of the lung/trachea/bronchus was 5.9 per 100 000, up from 5.1 in the previous reporting period. The incidence increased with increasing age, from 1.9 per 100 000 in those aged 35-39 years, up to 46.0 per 100 000 in those aged 65-69 years.




Figure 14: Age-specific Incidence rates per 100 000 in 2010-2014, Trachea, bronchus and lung (C33-34)

Cancer of the lung/trachea/bronchus was the second most frequent cancer in White males (10.6%), and in the top five cancers among Coloured (9.8%), Nama (8.4%) and Baster (6.0%) ethnic groups.

Cancer of the lung/trachea/bronchus was the second most common cancer in Kharas (11.6%) and among the five most common cancers in the Hardap (8.2%), Erongo (7.5%) and Otjozondjupa (6.0%) regions.

Other notable cancers in males

Cancer of the larynx appeared as the sixth most common cancer among males in Namibia with a total of 147 cases reported during the period 2010 to 2014 (3.1% of all male cancers; ASR 5.3 per 100 000). The incidence of laryngeal cancer increased with age and was highest in those aged 75-79 years, with an ASR of 49.4 per 100 000. Although laryngeal cancer ranked as the sixth most common cancer overall among males, it appeared among the five leading cancers among the Nama (7.7%), Damara (7.0%) and Baster (3.9%) ethnic groups.

Colon cancer was the third most common cancer among White males (9.5%) and fifth most common among Coloured males (4.9%). Colon cancer was the seventh most common cancer among male Namibians with a total of 129 cases and an ASR of 4.2 per 100 000.

Liver cancer was the eighth most common cancer among males in Namibia comprising 2.6% of male cancers with an ASR of 4.0 per 100 000. Liver cancer was among the top five cancers among the Kavango (5.3%), Coloured (4.9%) and Ovambo (2.9%) ethnic groups.

Among the Ovambo Kaposi sarcoma, prostate, eye, Non-Hodgkin lymphoma (ASR 3.1) and liver cancers were the five leading cancers in males, accounting for 28.6%, 24.1%, 6.1%, 4.8% and 2.9% of all cancers respectively.

Among White males, cancers of the prostate, lung/trachea/bronchus, colon, Non-Hodgkin lymphoma and melanoma of the skin were ranked the leading cancers, accounting for 19.9%, 10.6%, 9.5%, 6.0% and 5.0% of all male cancers respectively.

Leading cancers among females

The predominant cancer in women was breast cancer (27.3%, ASR 40.2), followed by cancers of the cervix (19.4%, ASR 27.2), Kaposi sarcoma (8.8%, ASR 10.1), eye (4.0%, ASR 4.8), ovary (3.0%, ASR 4.3), colon, Non-Hodgkin lymphoma (2.8%, ASR 3.4), corpus uteri (2.1%, ASR 3.3) and lung / trachea / bronchus (1.8%, ASR 2.9) cancers.³




Figure 15: Bar Chart showing Top 10 Cancers in Namibia (2010-2014), excluding C44, Female

³ The ranking of top cancers in the report is based on age standardized rates, while the ranking depicted in the bar chart is on absolute numbers.

Breast cancer

Breast cancer was the most common cancer among Namibian women between 2010 and 2014. A total of 1 579 cases were reported, comprising 27.4% of all female cancers. The ASR for female breast cancer was 40.2 per 100 000; a significant increase from 26.1 in the previous reporting period. The annual incidence increased with increasing age, peaking at 189.1 per 100 000 in women 70-74 years of age.




Figure 16: Age-specific Incidence rates per 100 000 in 2010-2014, Breast (C50)

Breast cancer was the most common cancer in most ethnic groups; White (44.7%), Coloured (38.3%), Nama (35.7%), Baster (31.4%), Damara (29.5%), Nama (28.7%), Herero (26.5%), Ovambo (24.5%), Kavango (21.5%), Tswana (17.6%) and Caprivi (11.8%) women.

Breast cancer was the most common cancer in most regions in Namibia, Omaheke (39.0%), Hardap (37.2%), Erongo (35.8%), Otjozondjupa (34.4%), Kharas (32.6%), Khomas (30.2%), Omasuti (25.7%), Oshana (24.9%) and Kavango (22.1%) regions, and the second most common cancer in Oshikoto (24.4%), Kunene (21.4%), Ohangwena (21.0%) and Caprivi (12.7%) regions.

Cervix cancer

Cervical cancer was the second most common cancer among women with 1 118 diagnoses (19.4% of all cancers in females) in the period beginning in 2010 and ending in 2014. The ASR for cervical cancer was 27.2 per 100 000 representing an almost two-fold increase since the previous reporting period when the ASR was 15.6 per 100 000. The annual incidence of cervical cancer increased with increasing age, from 0.2 per 100 000 in women aged 20-24 peaking at 122.8 per 100 000 in women aged 75-79 years.




Figure 17: Age-specific Incidence rates per 100 000 in 2010-2014, Cervix uteri (C53)

Cervical cancer was either the leading cancer or in the top three cancers in all ethnic groups, ranging from 4.8% of female cancers in White women to 33.1% among Caprivi women.

Cervical cancer was among the top two cancers in each region comprising between 9.9% of cancers in the Erongo Region and 38.7% of all cancers in the Caprivi Region.

Kaposi sarcoma

Kaposi sarcoma (KS) was the third most common cancer among women in this five-year reporting period. A total of 507 cases were reported, comprising 8.8% of all female cancers. The ASR was 10.1 per 100 000 up from 6.8 per 100 000 in the previous reporting period (2006-2009) and 4.1 in the prior reporting period (2000-2005). Annual incidence according to age showed a bell-shaped curve, increasing with age and peaking at 32.5 per 100 000 women among those who were aged 30 to 34 years and then decreasing in older women.




Figure 18: Age-specific Incidence rates per 100 000 in 2010-2014, Kaposi sarcoma (C46)

KS was in the top five most common female cancers in the Tswana (17.6%), Caprivial (13.2%), Ovambo (12.1%), Kavango (10.8%), Herero (5.1%), Baster (4.4%), Damara (3.2%) and Coloured (2.0%) ethnic groups.

It was also among the three most common cancers in all of the regions of Namibia, ranging between 4.5% (Erongo) and 14.5% (Kavango) of all cancers. In the Khomas and Kavango regions it ranked fourth most common cancer among Namibian women.

Eye cancer

Cancer of the eye was the fourth most common cancer among Namibian women during the period under review. A total of 232 cases were reported, comprising 4.0% of all female cancers. The ASR for ovarian cancer was 4.8 per 100 000 compared to an ASR of 1.3 in the previous reporting period. Annual incidence also showed an approximately bell-shaped curve with age, peaking at 12.3 per 100 000 in women aged 45-49 years.




Figure 19: Age-specific Incidence rates per 100 000 in 2010-2014, Eye (C69)

Eye cancer was among the third most common cancer among Herero (5.3%) and Damara (4.5%) women, and fourth most common cancer among Tswana (8.8%) and Ovambo (5.1%) women.

Regionally, eye cancer appeared among the five leading cancers in Oshana (7.6%), Khomas (6.0%), Kunene (5.4%) and White (3.3%) females.

Ovarian cancer

Cancer of the ovary was the fifth most common cancer among Namibian women during the period 2010 to 2014. A total of 175 cases were reported, comprising 3.0% of all female cancers. The ASR for ovarian cancer was 4.3 per 100 000, having increased from an ASR of 2.9 in the previous reporting period. Annual incidence increased with age, peaking at 20.1 per 100 000 in women aged 70-74 years.




Figure 20: Age-specific Incidence rates per 100 000 in 2010-2014, Ovary (C56)

Ovarian cancer was among the five most common cancers among Herero (5.1%), Coloured (4.8%), Damara (3.6%) and Capriviian (3.2%) women.

Regionally, ovarian cancer appeared among the five leading cancers in Otjozondjupa (6.6%), Kunene (5.4%), Omaheke (4.8%), Kharas (4.3%), Erongo (4.1%), Oshana (3.3%) and Oshikoto (2.7%) women.

Other notable cancers in females

Non-Hodgkin lymphoma was the sixth most common cancer among females between 2010 and 2014, comprising 2.8% of all cancers in women, ASR 3.4 per 100 000 – up from 2.3 in 2006-2009, and showed an increasing incidence according to age group. The peak incidence occurred in women aged 75-79 years of age (ASR 15.8 per 100 000).

Corpus uteri cancer was the seventh most common cancer among women overall (2.1% of all cancer in women, ASR 3.3 per 100 000 – up from 2.2 per 100 000 in 2006-2009) and showed an increasing incidence with increasing age, peaking in women aged 65-69 years old (ASR 28.4 per 100 000).

Cancer of the lung/trachea/bronchus was the eighth most common cancer among women accounting for 1.8% of all cancers and ASR of 2.9 per 100 000 women, compared to 2.3 per 100 000 in 2006-2009. Lung cancer was the fourth most common cancer among White and Coloured women accounting for 4.4 and 4.0% of all cancers in these women.

Melanoma of the skin was the fourth most common cancer among Kavango women comprising 5.5% of cancers. Melanoma of the skin was also the fifth most common cancer among White women, comprising 4.4% of all cancers.

Colon cancer was the second most common cancer among White women comprising 5.9% of cancers in this group. Cancer of the mouth was the fifth most common cancer among Ovambo women, comprising 1.3% of cancers.

Childhood cancers

A total of 357 cancer cases were recorded in children (defined as a person below the age of 15) during the five-year period from 2010 to 2014, of which 52.1% were male. Excluding other skin cancers, 353 cases were recorded among children (ages 0-14). The joint most common cancer in male children was kidney cancer and Non-Hodgkin lymphoma (27 cases in each, 14.8% of the total in each), followed by eye cancer (19 cases, 10.4%), brain cancer (16 cases, 8.7%), lymphoid leukemia (15 cases, 8.2%) and Kaposi sarcoma (13 cases, 7.1%).

Among female children there were 24 cases (14.2%) of eye cancer, 18 cases (10.7%) of kidney cancer, 17 cases (10.1%) of Non-Hodgkin lymphoma, 16 cases (9.5%) of brain cancer and 13 cases (7.7%) of Kaposi sarcoma. (These data are not tabulated, but the ASR's are included in Tables 1 and 2 and for each of the years separately in the appendix tables.)

Basis of diagnosis of cancers

The proportion of all cancers that were histologically verified between 2010 and 2014 was 94.8%. When (other skin) was excluded this dropped to 94.5%. This represents a substantial increase compared to the previous reporting period when these proportions were 86.4% and 82.6% respectively.

The cancer most likely to be diagnosed on death certificate only was Kaposi sarcoma: 16.1% in males and 15.4% in females. The most common cancers diagnosed by only clinical means were brain cancer (31.9% in males and 22.1% in females), pancreatic cancer (17.9% in males and 23.1% in females) and lung/trachea/bronchus cancers (21.1% in males and 16.2% in females). The data for basis of diagnosis are shown in Tables 7 and 8). Also see appendix for data quality indicators.

DISCUSSION

Limitations

The annual number of cancer cases reported between 2010 and 2014 appears to have stabilised between 1 900 and 2 600 cases. While this still shows substantial variability, there is less variability than in the previous reporting period when the annual number of cases varied between 1 100 and 2 800. This suggests that vigilance in standardising cancer registration procedures are paying dividends and the data reported here for 2010 to 2014 more likely reflect the true annual burden of cancer in Namibia. Stability in reported numbers and estimated ASRs is imperative for clear interpretation of cancer trends over time.

Increasing burden of cancer in Namibia

The significant increase in the overall incidence of cancer (excluding non-melanoma skin cancers) in both males and females in Namibia from 100.7 per 100 000 to 144.2 per 100 000 in males, and from 90.2 per 100 000 to 138.7 per 100 000 in females from the previous reporting period needs to be explored carefully. It is likely that the NNCR is now better collating all the cancer data in the country having been running the registry since 1995.

The tripling in the rate of eye cancer in females and a doubling of the rate of prostate, Non-Hodgkin lymphoma and eye cancer in males, and breast and cervical cancer and Kaposi sarcoma in females is a major concern.

Cancers associated with Human Immunodeficiency Virus infection

The prevalence of Human Immunodeficiency Virus (HIV) among pregnant women in Namibia reached its peak of approximately 22.0% in 2002 and has since appeared to steadily decline to 17.8% in 2008 and 16.9% in 2015 (Ministry of Health and Social Services Surveillance Report of the 2012 National HIV Sentinel Survey, Republic of Namibia, November 2014). The occurrence of HIV associated cancers has, as in many other high HIV prevalence African countries, increased over time. Declining incidence of HIV associated cancers is likely to lag behind declining HIV prevalence and incidence estimates, and the impact of improving access to HIV treatment on cancer patterns is yet to be noted in Namibia. Kaposi sarcoma dropped from most common to the second most common cancer among males and remained the third most common in females in 2010 to 2014 compared to 2006-2009. Kaposi sarcoma, as a proportion of all cancers, has shown small decreases in both sexes in the most recent reporting period, but in both cases the ASR has increased. Non-Hodgkin lymphoma was the fifth most common cancer in males and the sixth most common in females in 2006 to 2009, and in 2010 to 2014 was the third most common cancer in males and remained the sixth most common cancer in females, but the proportion of all cancers and the ASR that NHL represents has increased. Eye cancer increased its ranking and is now the fourth most common cancer among males and females in Namibia. The proportion of HIV associated cancers has unsurprisingly shown a trend of increase over time.

The appearance of cancers with an infectious aetiology, such as KS, NHL and eye cancers among the leading cancer sites, is undoubtedly linked to the HIV/AIDS epidemic. Since the advent of HIV, KS has become the leading cancer in men, and the second leading cancer in women in Zimbabwe and most central and east African countries. Those aged 15 to 44 (the most sexually active age group) are most affected. The human herpes virus 8 (HHV-8) / Kaposi sarcoma associated herpes virus (KSHV), is the necessary causal agent in the development of KS. The relationship between HIV and HHV8 in causing Kaposi sarcoma has now been well described. Other known risk factors for KS include increasing age and male sex.

The current incidence rates for KS in Namibia are still likely to be an underestimation of the true rates. Firstly, biopsies are usually only performed for patients with a single lesion or very early lesions, or if the doctor is inexperienced in recognising KS. Many KS patients are already so ill that they are treated supportively and no biopsies are performed. In addition, patients from peripheral areas with advanced illness are treated supportively and are unlikely to be referred to the sentinel areas. The registry would therefore not record these cases. It should however be noted that the histological verification of KS has increased substantially from below 40% in the previous reporting period to approximately 84% in the current reporting period, with concomitant declines in clinical diagnosis from approximately 60% to less than 1% in 2010-2014.

If the HIV epidemic progresses in Namibia as it has in other African countries, then the incidence of KS will increase dramatically in coming years. Other cancers that are likely to increase because of HIV immunosuppression include non-Hodgkin lymphoma and, possibly, conjunctival (eye) cancers (Newton et al. 2002). It is unclear whether cervical cancer and other female genital cancers, which also have a viral aetiology, will increase in incidence as a result of HIV. However, an HIV prevention program, if effectively implemented, should ultimately reduce the incidence of HIV-related cancers in men, women and children (International Collaborative Group on HIV and Cancer, 2000). Effective rollout of antiretroviral treatment for HIV will also likely reduce the incidence of HIV-associated cancers. The surveillance of cancer patterns in Namibia, particularly within different regions and ethnic groups, would form an essential part of such a program.

Prostate cancer

Prostate cancer has replaced Kaposi sarcoma as the most common cancer among males overall in the period 2010 to 2014, and the incidence has increased from 23.5 per 100 000 in the previous reporting period (2005-2009) to 37.6 per 100 000 in 2010 - 2014. Prostate cancer was among the five most frequent cancers in almost all ethnic groups and regions examined. Prostate cancer is one of the leading cancers in men worldwide. Incidence rates vary from 1 to 100 per 100 000 in different areas of the world, suggesting that regional factors play a role in its aetiology. Risk factors for prostate cancer include a diet high in fat and low in vegetables. There is suspicion that infections may play a role, but the agent(s) responsible have not been identified. In many western countries, the incidence of prostate cancer is increasing, possibly due to the increasing use of prostate specific antigen (PSA) as a screening tool. Further investigation is warranted to explore reasons for the recent increase in incidence of prostate cancer, and to determine risk factors for prostate cancer specific to Namibia.

Tobacco and alcohol related cancers

Cancers of the oral cavity, larynx and oesophagus were among the leading cancers in Namibian males and females. Internationally, tobacco smoking and alcohol consumption are the main risk factors for these cancers, with the two factors combined having a multiplicative effect on risk. Lung cancer was also among the five leading cancers in certain ethnic groups (Coloured, Baster and Nama). The main cause of lung cancer is tobacco smoking, for which there is a clear dose-response effect, related to both duration of smoking and amount smoked.

Tobacco is a risk factor for cancer both in its smoked and chewed form, and in the form of oral snuff. Chewing the areca (betel) nut is a dominant risk factor for oral cancer in certain communities worldwide, including in Southern Africa. Pipe-smoking is a risk factor for lip cancers. It is essential to monitor amounts and types of alcohol and tobacco consumption by Namibians with particular cancers, so that risk factors specific to this population can be determined, and preventative programs instituted.

Oral cancers and oesophageal cancer tended to decline during the period covered by this report. The declines may be due to as yet undefined changes in alcohol use or smoking habits. The male to female ratios for tobacco and alcohol related cancers in Namibia were 5.3, 3.0, 3.0 and 2.0 for cancers of the larynx, oesophagus, tongue and lung respectively. These ratios may reflect different patterns of tobacco and alcohol use among males and females, however further study is required to clarify the risk factors.

Additional risk factors exist for these cancers in different ethnic / regional groups, e.g. exposure to fungal toxins (oesophageal cancer), domestic and industrial pollution (larynx and lung) and ultra-violet radiation (lip). The relative importance of different risk factors for alcohol and tobacco related cancers needs to be determined for each population group in Namibia. Given the prominence of cancers of the upper aero digestive tract among certain ethnic groups, it would be important to investigate the existence of unusual risk factors for these cancers among Namibians.

Female-specific cancers

Breast cancer and cervical cancer were the two most common cancers among women in all ethnic and regional groups in Namibia. The ASR of both these cancers increased significantly in this reporting period relative to 2006 - 2009.

Risk factors for female breast cancer include early menarche, late age at first childbirth, a high-fat diet and certain genetic mutations, including BRCA1/2. Other possible risk factors include high alcohol consumption and the use of certain post-menopausal hormone replacement therapies. Public health campaigns that encourage women to present early to hospital for a breast exam could reduce mortality from this disease.

Risk factors for cervical cancer include infection with certain types of human papillomaviruses (HPV), early initiation of sexual intercourse, a history of multiple sexual partners (or a partner with multiple sexual partners), a history of sexually transmitted infections, cigarette smoking and long-term use of oral contraception. Lower socio-economic status has been associated with a higher risk of developing cervical cancer, possibly due to lack of access to good health care and Papanicolaou ('Pap-smear') tests. Although only a small proportion of women infected with HPV develop cervical cancer, recent studies have shown that women who have used hormonal contraceptives on a long-term basis (more than five years) are more likely to develop cancer of the cervix than those with less contraceptive use. HPV vaccination and annual cervical cancer screening programs have been demonstrated to reduce the incidence of this cancer, and it is recommended that such a program be implemented in Namibia on a national level by the Ministry of Health and Social Services. Cervical cancer rates among all race groups should be closely monitored as the HIV epidemic progresses. Furthermore, rollout of the HPV vaccine is likely to lead to significant reductions in cervical cancer.

Ovarian cancer was the fifth most common cancer among Namibian women. Risk factors for ovarian cancer include certain genetic mutations (BRCA1/2), delayed age at first pregnancy, lower parity, and a high-fat diet. Current users of oral contraceptives appear to be protected, although post-menopausal women who are on long-term hormone replacement therapy may be at increased risk. No cost-effective screening methods are currently available. As with the other cancers discussed, it would be important to obtain detailed information on these patterns of use that are specific to subgroups within Namibia that are associated with these female cancers.

Colon cancer

Colon cancer was the seventh most common cancer among Namibian men, and the tenth most common cancer among women. Risk factors for colorectal cancer include high-fat/low-fibre diets, inflammatory bowel disease, and a family history of colon cancer (particularly among first-degree relatives). High dietary folate has been shown to be protective. Rates of colorectal cancer are higher in western countries than in developing countries.

Basal and squamous cell skin cancers

For the purposes of this report, basal and squamous cell carcinomas of the skin (BCC and SCC; categorised as 'other skin') were excluded from rate calculations for 'all cancers', as well as frequency rankings of cancers. This is standard practice in cancer registry reports, thus this format allows international comparisons to be made. These lesions are often treated at outpatient facilities or doctors' rooms without histological confirmation, and rates are therefore underestimated. Despite these limitations, these non-melanoma skin cancers were the third most common cancer overall among males (27.1%, ASR 9.6 per 100 000) and the fourth most common among females (6.1%, ASR 7.6) in Namibia, a dramatic decrease in both sexes compared to the previous reporting period (2006-2009). Risk factors for these cancers include fair skin, albinism, and short, intense sun exposure. These cancers are therefore, as anticipated, most common among Whites. Given the high incidence and known risk factors, it is important to highlight these cancers in preventative and educational public health programs.

Cancers among children

It is uncertain whether the data used for this report were complete with respect to childhood cancers, despite the large increases noted compared to the previous reporting period. In addition to typical childhood cancers such as kidney cancer and eye cancer, HIV associated cancers such as eye, Non Hodgkin lymphoma and Kaposi sarcoma were among the top six cancers in both males and females aged less than 15 years. These together accounted for 32.3% and 32.0% of cancers among males and female children respectively.

RECOMMENDATIONS

This report has highlighted the leading cancers in Namibian males and females overall, and examined cancer frequencies in the different ethnic groups and geographic regions.

Calculation of incidence rates by gender, population group, ethnic group and region is important, as disease patterns differ due to differences in susceptibilities and exposures in different groups (e.g. genetic susceptibilities, socio-economic status, diet, alcohol and tobacco use). In addition, differences between groups help to identify inequalities in service delivery or utilisation. However, for this report, it was only possible to calculate incidence rates for the whole of Namibia (males and females), as population size estimates by 5-year age groups were not available for the different ethnic groups, or for regions.

The annual number of cases of cancer reported to the NNCR requires careful monitoring to examine peaks and troughs and to ensure that good quality cancer statistics can be provided and trends in cancer incidence be meaningfully interpreted.

Since these data are required for the development of cancer control program and advocacy, and to improve the cancer statistics for Namibia, it is recommended that cancer be made a reportable condition.

It is furthermore recommended that this report be used to guide the investigation of risk factors for the leading cancers in Namibia since key risk factor data is available on the NNCR database.

Accurate collection of regional data

During the analysis of the data it became apparent, as it had in the previous report, that cancer occurrence appeared to be much higher in Khomas than in other regions. It was ascertained that, if no address was stated on the histology report, the address was taken to be Windhoek, and coded as Khomas Region because of the treatment centre being in this city. We strongly recommend that in the future, missing data be recorded as such, and that efforts are made to accurately record region of residence. This would most likely be better achievable when cancer is made a reportable disease and full case data can be obtained.

If this is done and population denominators become available in 5-year age groups by sex, incidence rates can be calculated accurately for regions. Collecting such data correctly will in future highlight regional differences in cancers, which will help in determining risk factors and inform prevention program.

Provision of population statistics by ethnic group

Examining cancer incidence by ethnic group is extremely important in cancer epidemiology, as different ethnic groups have different genetic susceptibilities to cancer, and are exposed to different environmental and behavioural risk factors. They therefore display different cancer patterns, which require public health interventions tailored to specific ethnic groups.

Patterns such as these can only be accurately determined if age standardised incidence rates are calculated; however estimates of the numbers of people in each grouping are required for such calculations. It is therefore strongly recommended that such figures be obtained for future cancer registry reports and that the NNCR advocate the Namibian Central Bureau of Statistics to collect and provide these data in the next census. A minimal requirement would be population estimates for 'white', 'black' and 'mixed' ethnic groupings.

CONCLUSION

The NNCR has been successful in producing a data set that indicates the overall occurrence of cancer in Namibia. This data has highlighted the increasing burden of cancer in Namibia as well as the leading cancers, which should form a basis for research into risk factors for cancer specific to the Namibian population. Refinement of the data collection process and access to more detailed population statistics will hopefully allow more detailed reports to be produced in the future. Nevertheless, the current report indicates that the NNCR is an essential resource and an asset to the public health system.

REFERENCES

Carrara H, Stein L, Sitas F, Zietsman A, Gariseb B, Rautenbach N, Pontac J, Johannesson I and Ferlay J. Cancer in Namibia 2000-2005, Namibia Cancer Registry, Windhoek 2009.

Carrara H, Stein L, Sitas F, Zietsman A, Gariseb B, Rautenbach N, Pontac J, Rooi T, and Taapopi L. Cancer in Namibia 2006-2009, Namibia Cancer Registry, Windhoek 2011.

Cooke A, Parkin DM and Ferlay J. Canreg5 Software (c). IARC, Lyon 2001.

Cooke A, Parkin DM and Ferlay J, Descriptive Epidemiology © 2002-2005, International Agency for Research on Cancer, World Health Organisation.

Ferlay J, Burkhard C, Whelan S and Parkin DM. Check and Conversion Programs for Cancer Registries (IARC/IACR Tools for Cancer Registries). IARC Technical Report No.42 Lyon, 2005.

Ferlay J, Soerjomataram I, Ervik M, Dikshit R, Eser S, Mathers C, Rebelo M, Parkin DM, Forman D, Bray, F. GLOBOCAN 2012 v1.0, Cancer Incidence and Mortality Worldwide: IARC CancerBase No. 11 [Internet]. Lyon, France: International Agency for Research on Cancer; 2013. Available from: <http://globocan.iarc.fr>.

Fritz A, Percy C, Jack A, Shanmugaratnam K, Sabin L, Parkin DM and Whelan S (eds). International Classification of Diseases for Oncology, Third Edition. WHO, Geneva 2001.

International Collaboration on HIV and Cancer. Highly active antiretroviral therapy and the incidence of cancer in HIV infected adults. JNCI 2000; 92: 1823-1830.

Jensen OM, Parkin DM, MacLennan R, Muir CS, Skeet RG. 1991. Cancer Registration Principles and Methods: World Health Organisation, International Agency for Research on Cancer.

Koehler S. Namibia's cancer atlas: 1989-1994. Namibia National Cancer Registry 1997.

Ministry of Health and Social Services Surveillance Report of the 2012 National HIV Sentinel Survey, Republic of Namibia, November 2014.

Newton R, Sitas F, Dedicoat M, Ziegler JL. HIV infection and Cancer. Essex M, Mboup S, Kanki PJ, Marlink R, Tlou SD (eds). AIDS in Africa, 2nd Edition. Raven Press 2002.

Parkin DM, Whelan SL, Ferlay J, Raymond L and Young J. Cancer Incidence in five continents, Vol VII, IARC Scientific Publication 143, IARC, Lyon 1997.

Parkin DM, Ferlay J, Hamdi-Chérif M. Sitas F, Thomas JO, Wabinga H, Whelan SL. Cancer in Africa. Epidemiology and Prevention. IARC Scientific Publications No.153. IARC Press, Lyon 2003.

Parkin DM, Sitas F, Chirenje M, Stein L, Abratt R, Wabinga H. Part I: Cancer in Indigenous Africans--burden, distribution, and trends. Lancet Oncol. 2008 Jul;9(7):683-92. Review.

Sitas F, Parkin DM, Chirenje M, Stein L, Abratt R, Wabinga H. Part II: Cancer in Indigenous Africans--causes and control. Lancet Oncol. 2008 Aug;9(8):786-95. Review.

The Republic of Namibia Central Bureau of Statistics National Planning Commission. Population Projections 1991-2021 Namibia National and Regional Figures, Namibia 2001.

Wabinga HR, Parkin DM, Wabwire-Mangen F, Mugerwa JW. Cancer in Kampala, Uganda in 1989-1991: Changes in Incidence in the era of AIDS. Int. J. Cancer 1993; 54: 23-26.

World Health Organisation. National Cancer Control Program. Policies and managerial guidelines. 2nd edition. Geneva 2002.

The following summary measures have been used in tables:

- *Site* – a shortened version of the full ICD-10 title describes each site or site grouping
- *All ages* – the total number of cases by site and for all sites
- *Age unk* – the number of cases of unknown age by site
- *0-, 5-, 10-...* – the age groups for which incidence data is presented
- *Crude rate* – the crude average annual incidence rate, based on the total number of registrations by site
- *%* - The proportional frequency of each site to the total of all sites excluding ICD10 C44 (Other skin)
- *CUM 0-64* – the sum over each year of age of the age-specific incidence rates, taken from birth to age 64
- *CUM 0-74* – the sum over each year of age of the age-specific incidence rates, taken from birth to age 74
- *ASR(W)* – the world age-standardized incidence rate by site
- *ICD(10th)* – the ICD-10 code(s) corresponding to the site or group of sites given in the left hand column

Table 1: NAMBIA (2010-2014) Annual incidence per 100 000 by age group - Male

SITE	ALL AGE AGES UNK	Central Cancer Registry, NAMIBIA (2010-2014)																CRUDE RATE (%)	CUM CUM ASR ICD (10th)			
		0-	5-	10-	15-	20-	25-	30-	35-	40-	45-	50-	55-	60-	65-	70-	75-	80+				
Lip	6 0	-	-	-	0.2	0.2	-	-	0.6	-	1.2	3.6	19.7	12.0	17.3	11.9	25.6	14.5	15.6	0.1 0.1 0.00 0.01 0.1 C00		
Tongue	103 0	-	-	-	0.2	-	0.2	0.9	0.8	0.3	-	6.8	9.1	12.9	14.1	25.2	21.7	29.1	26.0	2.0 2.2 0.27 0.46 3.9 C01-02		
Mouth	116 0	-	-	-	0.2	-	0.2	0.4	0.5	1.0	1.7	3.6	2.1	4.6	2.2	3.0	5.9	5.8	5.2	2.2 2.4 0.23 0.46 3.9 C03-06		
Salivary glands	41 0	-	-	0.2	0.2	-	0.2	0.4	0.5	-	-	1.0	2.8	1.8	4.3	1.5	7.9	5.8	1.7	0.8 0.9 0.08 0.13 1.2 C07-08		
Tonsil	24 0	-	-	-	-	0.2	-	0.4	-	0.3	-	-	0.7	1.8	1.1	3.0	2.0	8.7	-	0.5 0.5 0.05 0.10 0.8 C09		
Other oropharynx	11 0	-	-	-	-	-	-	-	-	0.3	-	-	0.7	1.8	1.1	4.4	-	-	3.5	0.2 0.2 0.02 0.04 0.4 C10		
Nasopharynx	23 0	-	-	0.2	0.2	0.2	0.4	0.4	-	0.3	0.4	2.1	1.4	1.8	1.1	4.4	-	-	3.5	0.4 0.5 0.04 0.06 0.6 C11		
Hypopharynx	6 0	-	-	-	-	-	-	-	-	-	-	0.7	2.8	1.1	-	-	2.9	-	-	0.1 0.1 0.02 0.02 0.2 C12-13		
Pharynx unspecified	5 0	-	-	-	-	-	-	-	0.3	-	-	0.7	-	-	1.1	-	3.9	-	-	0.1 0.1 0.01 0.03 0.2 C14		
Oesophagus	108 0	-	0.2	-	-	-	-	0.5	0.6	0.8	3.6	9.8	12.9	21.7	28.2	25.6	8.7	19.1	2.1 2.3 0.25 0.52 3.9 C15			
Stomach	72 0	-	-	-	-	-	-	0.4	0.3	0.6	2.1	4.2	6.3	5.5	2.2	14.8	13.8	40.7	10.4	1.4 1.5 0.18 0.25 2.3 C16		
Small intestine	6 0	-	-	-	-	-	-	-	-	-	-	0.5	0.9	1.1	1.0	-	-	-	2.3	0.4 0.4 0.01 0.01 0.2 C17		
Colon	129 0	-	-	-	-	-	0.2	0.9	1.9	1.9	3.7	8.3	16.2	12.0	11.9	26.7	11.8	34.9	5.2	2.5 2.7 0.29 0.48 4.2 C18		
Rectum	69 0	0.1	-	-	-	-	0.4	0.4	0.5	2.2	2.5	3.6	3.5	2.8	6.5	11.9	15.8	20.3	8.7	1.3 1.4 0.11 0.25 2.1 C19-20		
Anus	23 0	-	-	-	-	0.3	-	-	-	0.2	-	0.8	2.1	4.9	-	-	1.5	5.9	5.2	0.4 0.5 0.04 0.08 0.7 C21		
Liver	124 0	0.3	-	0.2	0.2	0.2	0.4	0.4	-	0.3	0.4	2.1	1.4	1.8	1.1	4.4	-	-	3.5	0.4 0.5 0.04 0.06 0.6 C22		
Gallbladder etc.	12 0	-	-	-	-	-	-	-	-	-	-	0.5	0.7	2.2	3.0	2.0	-	-	8.7	0.2 0.3 0.02 0.04 0.4 C23-24		
Pancreas	28 0	-	-	-	-	-	-	-	-	-	-	3.5	3.7	7.6	7.4	3.9	5.8	1.7	1.7	0.6 0.08 0.13 1.0 C25		
Nose, sinuses etc.	20 0	0.1	-	-	-	-	-	0.6	0.2	-	-	0.5	2.1	0.9	4.3	3.0	-	-	5.8	0.4 0.4 0.05 0.06 0.6 C30-31		
Larynx	147 0	-	-	-	-	-	-	-	0.7	-	-	0.8	4.2	9.8	17.5	23.9	44.5	37.4	49.4	22.5	2.8 3.1 0.28 0.69 5.3 C32	
Trachea, bronchus and lung	161 0	-	-	-	-	-	-	-	-	1.9	3.3	5.2	5.6	36.0	24.9	46.0	41.4	17.4	15.6	3.1 3.4 0.39 0.82 5.9 C33-34		
Other thoracic organs	8 0	0.3	-	-	-	-	-	-	0.2	-	-	0.5	-	1.8	-	-	-	-	0.2 0.2 0.02 0.02 0.2 C37-38			
Bone	72 0	-	-	0.3	0.7	1.7	1.5	1.6	1.3	1.0	1.7	1.0	2.8	3.7	5.4	4.4	5.9	-	13.9	1.4 1.5 0.11 0.17 1.7 C40-41		
Melanoma of skin	88 0	-	-	-	-	-	-	-	0.2	1.9	1.6	1.7	3.1	8.4	7.4	6.5	13.3	9.9	23.3	29.5	1.7 1.8 0.15 0.27 2.6 C43	
Other skin	338 0	-	-	-	0.5	1.2	2.8	2.5	9.4	8.3	7.0	17.7	17.6	14.8	26.0	41.5	65.0	81.4	62.4	6.5 7.1 0.54 1.07 9.6 C44		
Mesothelioma	7 0	-	-	-	-	-	-	-	-	-	-	0.4	0.5	-	-	2.2	1.5	2.0	2.9	0.1 0.1 0.02 0.03 0.3 C45		
Kaposi sarcoma	882 0	0.4	0.8	0.8	2.0	1.7	9.4	48.9	49.1	62.4	54.7	50.6	30.5	39.0	31.1	39.4	46.5	29.5	16.9 18.4 1.75 2.10 2.2 C46			
Connective and soft tissue	91 0	0.8	0.8	-	1.4	1.5	0.9	4.3	1.3	1.7	1.0	4.9	4.6	8.7	5.9	3.9	14.5	5.2	1.7 1.9 0.16 0.21 2.2 C47,C49			
Breast	62 0	-	-	-	-	-	-	-	0.4	1.6	1.3	1.2	3.6	4.9	7.4	3.3	10.4	7.9	11.6	12.1	1.2 1.3 0.12 0.21 1.9 C50	
Penis	52 0	-	-	-	-	-	-	-	0.2	0.8	1.6	1.7	4.7	4.9	3.7	1.1	7.4	-	-	20.3	10.4	1.0 1.1 0.09 0.13 1.5 C60
Prostate	1128 3	-	-	-	-	-	-	-	-	1.6	0.6	1.7	7.3	24.6	64.6	147.4	289.2	425.7	546.5	449.0	21.6 23.6 1.24 4.83 3.7 C61	
Testis	32 0	0.4	0.2	-	-	0.3	0.4	0.9	1.3	0.3	1.7	1.6	1.4	0.9	-	-	1.5	-	-	0.6 0.7 0.05 0.05 0.7 C62		
Other male genital organs	10 0	0.1	-	-	-	0.2	0.2	-	-	-	-	1.4	0.9	-	-	-	-	-	2.0	0.2 0.2 0.02 0.03 0.3 C63		
Kidney	88 0	2.1	1.1	0.7	0.5	-	-	0.3	0.3	3.3	3.1	3.5	11.1	5.4	14.8	5.9	14.5	3.5	1.7 1.8 0.16 0.26 2.5 C64			
Renal pelvis	1 0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.0 0.0 0.00 0.00 0.0 C65			
Uterus	0 0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.0 0.0 0.00 0.00 0.0 C66			
Bladder	72 0	0.1	0.3	0.2	-	0.2	-	0.3	1.0	0.8	1.0	3.5	6.5	13.0	22.2	17.7	11.6	12.1	1.4 1.5 0.13 0.33 2.5 C67			
Other urinary organs	2 0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.0 0.0 0.00 0.00 0.1 C68			
Eye	241 0	0.2	0.2	-	0.8	0.7	1.8	7.0	15.1	16.5	15.6	7.0	14.8	8.7	19.3	13.8	14.5	5.2	4.6 5.0 0.45 0.62 6.2 C69			
Brain, nervous system	69 0	0.7	0.5	1.3	1.0	0.7	0.7	1.9	0.6	1.7	2.6	2.8	5.5	3.3	4.4	5.9	2.9	3.5	1.0 1.4 0.12 0.17 1.7 C70-72			
Thyroid	21 0	-	-	-	-	-	-	0.4	0.3	0.6	1.2	1.0	1.7	4.6	2.2	1.5	-	-	0.4 0.4 0.06 0.07 0.6 C73			
Adrenal gland	3 0	0.1	0.2	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.1 0.1 0.01 0.01 0.1 C74			
Other endocrine	5 0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.1 0.1 0.02 0.02 0.1 C75			
Hodgkin disease	32 0	0.1	0.6	0.3	0.7	0.4	0.7	0.5	1.3	1.7	0.7	2.8	-	-	1.5	-	-	-	0.6 0.7 0.05 0.06 0.7 C81			
Non-Hodgkin lymphoma	231 0	-	1.8	0.8	1.0	0.6	3.4	4.0	7.1	9.5	16.2	16.2	13.8	15.2	14.8	23.6	23.3	12.1	-	4.4 4.8 0.43 0.65 6.3 C82-85,C96		
Immunoproliferative diseases	2 0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.0 0.0 0.00 0.00 0.1 C88			
Lymphoid leukaemia	24 0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.5 0.5 0.05 0.11 0.8 C90			
Myeloid leukaemia	51 0	0.7	1.3	0.3	0.7	-	0.4	0.9	0.5	1.3	1.7	2.7	3.4	3.4	3.5	-	-	-	1.0 1.1 0.07 0.12 1.3 C91			
Leukaemia unspecified	2 0	0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1.0 1.1 0.09 0.15 1.4 C92-94			
Myeloproliferative disorders	8 0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.1 0.1 0.01 0.01 0.1 MDS			
Myelodysplastic syndromes	0 0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.1 0.1 0.02 0.02 0.2 MDS			
Other and unspecified	213 1	0.7	-	0.3	0.3	0.4	1.3	0.8	3.5	4.5	10.9	12.6	24.0	28.2	41.5	33.5	43.6	19.1	4.1 4.4 0.45 0.83 6.9 O&U			
All sites	5125 4	11.5	8.9	7.3	13.5	14.6	32.9	95.6	110.4	146.0	207.9	289.5	380.4	489.0	806.9	930.1	1162.8	842.5	98.0	9.04 17.74 153.8 ALL		
All sites but C44	4787 4	11.5	8.9	6.8	12.4	11.8	30.5	86.2	102.1	138.9	190.2	272.0	365.6	462.9	765.4	865.1	1081.4	780.1	91.5	100.0 8.51 16.66 144.2 ALLbc44		

, Reference population: World Standard Population

Table builtFri Nov 18 07:55:00 SAST 2016 by CanReg5.

Table 2: NAMIBIA (2010-2014) Annual incidence per 100 000 by age group – Female

Central Cancer Registry, NAMIBIA (2010-2014)

2012 Population Namibia

Incidence per 100,000 by age group (Period) - Female

SITE	ALL AGE AGES UNK
------	------------------

Table 3: NAMIBIA (2010-2014) Cancer frequencies by ethnic group – Males

SITE	White	Baster	%	San	Capriviian	Damara	Herero	Kavango	Coloured	Nama	Tswana	Ovambo	Other/Unk	n	%											
Up	1	0.2	1	0.4	0	0	0	1	0.3	0	1	0.6	0	0	0.1	0										
Tongue	11	2.4	6	2.1	4	23.5	0	0	22	5.3	18	4.5	5	2.4	3	1.8	8	5.2	0	0	27	1.2	5	1.5		
Mouth	8	1.7	8	2.8	2	2.5	22	5.3	13	3.3	2	1	10	6.1	14	9	3	7.5	36	1.6	10	3				
Salivary glands	4	0.9	2	0.7	1	5.9	0	0	2	0.5	6	1.5	1	0.5	1	0.6	3	1.9	0	0	14	0.6	8	2.4		
Tonsil	3	0.6	1	0.4	0	0	0	1	1.2	0	7	1.7	4	1	2	1	2	1.2	3	1.9	0	0	4	0.2	0	0
Other oropharynx	0	0	1	0.4	0	0	0	1	1.2	0	2	0.5	0	0	0	1	0.6	2	1.3	0	0	2	0.1	2	0.6	
Nasopharynx	1	0.2	3	1.1	1	5.9	1	2.1	4	1	2	0.5	2	1	2	1	2	1.3	0	0	2	0.1	3	0.9		
Hypopharynx	0	0	0	0	0	0	0	0	0	1	0.3	0	0	0	0	0	0	0	0	0	0	2	0.1	1	0.3	
Pharynx unspecified	0	0	1	0.4	0	0	0	0	0	0	0	0	0	0	1	0.5	0	0	1	0.6	0	0	2	0.1	0	0
Oesophagus	10	2.2	4	1.4	1	5.9	0	0	17	4.1	14	3.5	10	4.8	2	1.2	9	5.8	1	2.5	33	1.5	7	2.1		
Stomach	10	2.2	3	1.1	0	0	3	3.7	5	1.2	6	1.5	6	2.9	4	2.5	4	2.6	0	0	20	0.9	11	3.3		
Small intestine	0	0	0	0	0	0	1	1.2	2	0.5	1	0.3	1	0.5	0	0	0	0	0	1	0	0	0	0	0	0
Colon	44	9.5	11	3.9	0	0	2	2.5	6	1.5	13	3.3	3	1.4	8	4.9	1	0.6	1	2.5	34	1.5	6	1.8		
Rectum	16	3.5	7	2.5	0	0	1	1.2	3	0.7	4	1	4	1.9	2	1.2	0	0	0	29	1.3	3	0.9			
Anus	4	0.9	1	0.4	0	0	0	0	2	0.5	1	0.5	1	0.6	1	0.6	0	0	0	10	0.4	1	0.3			
Liver	9	1.9	2	0.7	0	0	3	3.7	8	1.9	7	1.8	11	5.3	8	4.9	4	2.6	1	2.5	65	2.9	6	1.8		
Gallbladder etc.	1	0.2	2	0.7	0	0	0	0	0	0	0	0	0	0	1	0.6	0	0	0	8	0.4	0	0			
Pancreas	9	1.9	4	1.4	0	0	0	0	4	1	2	0.5	1	0.5	5	3.1	0	0	0	3	0.1	0	0			
Nose, sinuses etc.	2	0.4	0	0	0	0	0	0	1	0.2	3	0.8	1	0.5	1	0.6	1	0.6	0	9	0.4	2	0.6			
Larynx	10	2.2	11	3.9	0	0	3	3.7	29	7	17	4.3	1	0.5	5	3.1	12	7.7	0	0	48	2.1	11	3.3		
Trachea, bronchus and lung	49	10.6	17	6	0	0	0	0	14	3.4	19	4.8	2	1	16	9.8	13	8.4	1	2.5	19	0.8	11	3.3		
Other thoracic organs	0	0	2	0.7	0	0	0	0	0	0	0	0	0	0	2	1.2	0	0	1	2.5	2	0.1	1	0.3		
Bone	3	0.6	7	2.5	0	0	0	0	0	0	8	1.9	5	0.8	3	1.4	2	1.2	0	0	4	1.1	7	2.1		
Melanoma of skin	23	5	0	0	0	0	0	1	0.2	6	1.5	5	2.4	1	0.6	0	0	1	2.5	47	2.1	4	1.2			
Other skin	73	15.8	29	10.2	2	11.8	3	3.7	13	3.1	13	3.3	14	6.7	12	7.4	9	5.8	0	0	148	6.6	22	6.5		
Mesothelioma	1	0.2	0	0	0	0	1	1.2	0	0	1	0.3	0	0	0	0	1	0.6	0	0	3	0.1	0	0		
Kaposi sarcoma	9	1.9	7	2.5	0	0	23	20.4	32	7.7	23	5.8	62	29.8	5	3.1	10	6.5	5	12.5	643	28.6	63	18.6		
Connective and soft tissue	6	1.3	6	2.1	0	0	0	0	0	1	0.2	5	1.3	1	0.5	0	0	0	0	51	2.3	7	2.1			
Breast	9	1.9	1	0.4	0	0	0	0	4	1	1	0.3	2	1	0	0	0	0	0	32	1.4	7	2.1			
Penis	4	0.9	4	1.4	1	5.9	4	4.9	5	1.2	2	0.5	2	1	2	1.2	4	2.6	1	2.5	23	1	0	0		
Prostate	92	19.9	96	33.9	2	11.8	16	19.8	97	23.5	116	29.1	21	10.1	36	22.1	24	15.5	14	35	543	24.1	72	21.3		
Testis	12	2.6	3	1.1	0	0	1	1.2	2	0.5	5	1.3	0	0	0	0	1	0.6	0	0	5	0.2	3	0.9		
Other male genital organs	0	0	1	0.4	0	0	0	0	0	3	0.7	1	0.3	2	1	0	0	0	0	2	0.1	1	0.3			
Kidney	10	2.2	7	2.5	0	0	0	0	6	1.5	11	2.8	8	3.8	3	1.8	5	3.2	2	5	29	1.3	7	2.1		
Renal pelvis	1	0.2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
Ureter	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
Bladder	13	2.8	17	6	0	0	0	0	0	4	1	7	1.8	4	1.9	2	1.2	2	1.3	0	0	19	0.8	4	1.2	
Other urinary organs	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2	0.1	0	0		
Eye	4	0.9	15	5.3	0	0	3	3.7	30	7.3	23	5.8	9	4.3	3	1.8	0	0	0	0	138	6.1	16	4.7		
Brain, nervous system	4	0.9	0	0	0	0	3	3.7	8	1.9	10	2.5	3	1.4	4	2.5	3	1.9	1	2.5	29	1.3	4	1.2		
Thyroid	1	0.2	2	0.7	0	0	0	0	0	1	0.3	2	1	0	0	0	0	0	0	9	0.4	4	1.2			
Adrenal gland	1	0.2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0			
Other endocrine	0	0	1	0.4	0	0	0	0	1	0.2	0	0	0	0	0	0	0	0	0	0	0	0	0			
Hodgkin disease	3	0.6	4	1.4	0	0	1	1.2	3	0.7	3	0.8	1	0.5	3	1.8	0	0	1	2.5	10	0.4	3	0.9		
Non-Hodgkin lymphoma	28	6	7	2.5	1	5.9	6	7.4	15	3.6	22	5.5	10	4.8	5	3.1	3	1.9	1	2.5	108	4.8	25	7.4		
Immunoproliferative diseases	1	0.2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
Multiple myeloma	5	1.1	1	0.4	0	0	0	1	1.2	6	1.5	7	1.8	0	0	0	0	0	0	0	1	0	0	0		
Lymphoid leukaemia	10	2.2	1	0.4	0	0	1	1.2	6	1.5	12	2.7	7	1.7	1	0.4	2	0.8	9	3.9	2	5.9	38	1.3		
Myeloid leukaemia	6	1.3	1	0.4	0	0	0	0	0	1.2	5	1	0.2	0	0	0	0	0	0	0	13	0.5	2	0.5		
Leukemia unspecified	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
Myeloproliferative disorders	2	0.4	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
Mylodysplastic syndromes	2	0.4	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
Other and unspecified	21	4.5	15	5.3	4	23.5	2	2.5	22	5.3	11	2.8	9	4.3	6	3.7	13	8.4	1	2.5	90	4	19	5.6		
All sites	536	312	19	84	426	426	222	175	164	40	2400	360	0	0	0	0	0	0	0	0	0	0	0	0		
All sites but C44	463	283	17	81	413	399	208	163	155	40	2252	358	0	0	0	0	0	0	0	0	0	0	0	0		

Table 4: NAMIBIA (2010-2014) Cancer frequencies by ethnic group – Females

SITE	White	Baster	%	San	Capriviian	Damara	Herero	Kavango	Coloured	Nama	Tswana	Ovambo	Other/Unk	n	%									
Up	0	0.0	1	0.0	0	0.0	0	0	1	0.2	0	0.0	0	4	0.1	0	0	0	0	0	0	0	0	
Tongue	7	1.5	3	1.1	2	10.0	0	0	10	2.3	3	0.7	0	0	4	1.6	6	2.6	0	0	12	0.4	1	0.3
Mouth	6	1.3	1	0.4	2	0	0	2	1.5	12	2.7	7	1.7	1	0.4	2	0.8	9	3.9	38	1.3	8	2.0	
Salivary glands	2	0.4	1	0.4	0	0	0	2	1.5	1	0.2	3	0.7	1	0.4	0	0	2	1.2	13	0.5	2	0.5	
Tonsil	0	0.0	2	0.7	0	0	0	0	0															

Table 5: NAMIBIA (2010-2014) Cancer frequencies by region – Males

SITE	Caprivi	Erlango	Hardap	Karas	Khomas	Kunene	Oshana	Kavango	Omahahe	Oshikoto	Otjozondjupa	Omasuti	Ohangwena	Unknown/Other
	n	(%)	n	(%)	n	(%)	n	(%)	n	(%)	n	(%)	n	(%)
Lip	0	0	0	0	1	0.5	1	0.6	3	0.2	0	0	0	0
Tongue	0	0	10	2.8	6	3.3	11	6.4	34	2.1	8	4.7	5	2.2
Mouth	2	2.6	16	4.4	12	6.5	8	4.7	30	1.8	8	7.4	15	1.5
Salivary glands	0	0	2	0.6	1	0.5	3	1.7	19	1.2	1	0.9	7	0.7
Tonsil	0	0	5	1.4	4	2.2	2	1.2	4	0.2	1	0.9	0	0
Other oropharynx	0	0	0	0	1	0.5	0	0	4	0.2	1	0.9	2	0.2
Nasopharynx	1	1.3	3	0.8	3	1.6	0	0	10	0.6	0	0	0	0
Hypopharynx	0	0	0	0	2	1.1	0	0	3	0.2	0	0	0	0
Pharynx unspecified	0	0	0	0	0	0	1	0.6	2	0.1	0	0	0	0
Oesophagus	1	1.3	10	2.8	4	2.2	7	4.1	40	2.5	5	4.6	9	0.9
Stomach	1	1.3	9	2.5	3	1.6	3	1.7	26	1.6	1	0.9	7	0.7
Small intestine	1	1.3	0	0	0	0	0	0	1	0.1	0	0	0	0
Colon	1	1.3	18	5	3	1.6	4	2.3	61	3.7	4	3.7	12	1.2
Rectum	0	0	7	1.9	4	2.2	0	0	30	1.8	0	0	13	3
Anus	1	1.3	1	0.3	2	1.1	2	1.2	3	0.2	0	0	3	1
Liver	2	2.6	19	5.3	5	2.7	2	1.2	28	1.7	0	0	27	10
Gallbladder etc.	0	0	0	0	1	0.5	1	0.6	4	0.2	0	0	3	0.3
Pancreas	0	0	6	1.7	4	2.2	2	1.2	10	0.6	0	1	0.1	0
Nose, sinuses etc.	2	2.6	2	0.6	1	0.5	0	0	6	0.4	1	0.9	2	0.2
Larynx	3	3.9	9	2.5	9	4.9	13	7.6	53	3.3	2	1.9	15	1.5
Trachea, bronchus and lung	0	0	27	7.5	15	8.2	20	11.6	58	3.6	4	3.7	2	0.2
Other thoracic organs	0	0	1	0.3	1	0.5	0	0	5	0.3	0	0	1	0
Bone	0	0	3	0.8	5	2.7	0	0	22	1.4	0	0	18	1.9
Melanoma of skin	1	1.3	8	2.2	0	0	2	1.2	16	1	3	2.8	22	2.3
Other skin	5	6.6	50	13.9	24	13	19	11	99	6.1	4	3.7	64	6.6
Mesothelioma	1	1.3	3	0.8	0	0	0	0	1	0.1	0	0	0	0
Kaposi sarcoma	24	31.6	51	14.2	18	9.8	16	9.3	181	11.1	7	6.5	240	24.7
Connective and soft tissue	1	1.3	9	2.5	4	2.2	5	1.9	27	1.7	2	1.9	20	2.1
Breast	0	0	7	1.9	2	1.1	1	0.5	23	1.4	0	0	15	1.5
Penis	0	3.9	4	1.1	1	0.5	2	1.2	18	1.1	0	0	14	1.0
Prostate	14	18.4	40	11.1	35	19	31	18	512	31.4	28	25.9	313	32.2
Testis	2	2.6	4	1.1	0	0	0	0	16	0	0	1	0.1	0
Other male genital organs	1	1.3	0	0	0	0	1	0.6	1	0.1	2	1.9	1	0.1
Kidney	0	0	13	3.6	6	3.3	2	1.2	30	1.8	1	0.9	3	0.3
Renal pelvis	0	0	0	0	0	0	1	0.6	0	0	0	0	0	0
Uterer	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Bladder	0	0	1	0.3	3	1.6	2	1.2	48	2.9	1	0.9	3	0.3
Other urinary organs	0	0	1	0.3	0	0	0	0	1	0.1	0	0	0	0
Eye	2	2.6	9	2.5	1	0.5	1	0.6	109	6.7	6	5.6	80	8.2
Brain, nervous system	3	3.9	8	2.2	2	1.1	0	0.5	109	4.3	3	2	4	3
Thyroid	0	0	1	0.3	2	1.1	0	0	6	0.4	0	0	7	0.7
Adrenal gland	0	0	1	0.3	0	0	0	0	2	0.1	0	0	0	0
Other endocrine	0	0	0	0	1	0.5	0	0	1	0.1	0	0	0	0
Hodgkin disease	0	0	6	1.7	2	1.1	0	0	6	0.4	0	0	7	0.7
Non-Hodgkin lymphoma	3	3.9	15	4.2	7	3.8	7	4.1	62	3.8	6	5.6	47	4.9
Immunoproliferative diseases	0	0	0	0	0	0	0	0	1	0.1	0	0	0	0
Multiple myeloma	0	0	4	1.1	0	0	1	0.6	7	0.7	1	0.9	1	0.2
Lymphoid leukaemia	1	1.3	7	1.9	3	1.1	1	0.6	13	7.0	3	2.8	21	2.2
Myeloid leukaemia	1	1.3	6	1.7	3	1.6	3	1.7	14	0.9	3	2.8	4	3
Leukaemia unspecified	0	0	0	0	0	0	1	0.6	0	0	0	0	0	0
Myeloproliferative disorders	0	0	0	0	0	0	0	0	3	0.2	0	0	0	0
Myelodysplastic syndromes	0	0	1	0.3	0	0	0	0	0	0.1	0	0	0	0
Other and unspecified	4	5.3	13	3.6	9	4.9	11	6.4	57	3.5	9	8.3	37	3.8
All sites	81	410	208	191	1727	112	1035	247	155	143	143	306	144	195
All sites but C44	76	360	184	172	1628	108	971	230	148	134	134	283	134	182

Table 6: NAMIBIA (2010-2014) Cancer frequencies by region – Females

SITE	Caprivi	Erlango	Hardap	Karas	Khomas	Kunene	Oshana	Kavango	Omahahe	Oshikoto	Otjozondjupa	Omasuti	Ohangwena	Unknown/Other
	n	(%)	n	(%)	n	(%)	n	(%)	n	(%)	n	(%)	n	(%)
Lip	0	0.0	0	0.0	1	0.4	1	0.1	0	0.3	0.2	0	0.0	0.0
Tongue	0	0.0	5	1.4	3	1.4	2	0.9	17	1.1	2	1.8	5	1.4
Mouth	3	2.0	5	1.4	3	1.4	5	2.1	22	1.4	5	4.5	13	1.0
Salivary glands	2	1.3	0	0.0	1	0.5	1	0.4	6	0.4	0	0.3	0	0.0
Tonsil	0	0.0	0	0.0	1	0.5	1	0.4	8	0.5	1	0.9	6	0.5
Other oropharynx	0	0.0	0	0.0	2	0.9	1	0.5	7	0.3	0	0.0	0	0.0
Nasopharynx	0	0.0	1	0.3	2	0.9	0	0.4	0	0.0	0	0.0	0	0.0
Hypopharynx	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Pharynx unspecified	0	0.0	0	0.0	0	0.0	0	0.1	1	0.0	0	0.0	0	0.0
Oesophagus	1	0.7	2	0.6	1	0.5	7	3.0	12	0.8	0	0.0	16	1.3
Stomach	0	0.0	7	1.9	2	0.9	23	1.4	0	0.0	13	1.0	4	1.4
Small intestine	0	0.0	0	0.0	1	0.5	0	0.2	0	0.1	2	0.7	0	0.0
Colon	0	0.0	14	3.9	2	0.9	6	2.6	45	2.8	0	0.0	10	0.8
Rectum	2	1.3	11	3.0	3	1.4	5	2.1	23	1.4	0	0	10	0.8
Anus	2	1.3	0	0.0	2	0.9	0	0.8	5	0.5	0	0	2	0.2
Liver	3	2.0	3	0.8	5	2.3	2	0.9	18	1.1	2	1.8	20	1.6
Gallbladder etc.	0	0.0	4	1.1	0	0.0	0	0.0	7	0.4	0	0.0	5	0.4
Pancreas	0	0.0	2	0.6	5	2.3	1	0.4	7	0.4	0	0.0	3	0.8
Nose, sinuses etc.	0	0.0	1	0.3	1	0.5	0	0.0	5	0.3	1	0.9	7	0.7
Larynx	0	0.0	3	0.8	1	0.5	0	0.0	20	1.3	0	0.0	7	0.0
Trachea, bronchus and lung	0	0.0	16	4.4	4	1.9	6	2.6	39	2.5	1	0.9	25	3.0
Other thoracic organs	0	0.0	0	0.0	0	0.0	0	0.0	1	0.1	0	0	0	0.0
Bone	1	0.7	4	1.1	0	0.0	1	0.4	21	1.3	3	2.7	1	0.7
Melanoma of skin	2	1.3	10	2.8	1	0.5	1	0.4	27	1.4	2	3.6	34	2.3
Other skin	4	2.7	28	7.7	25	11.6	25	10.7	100	6.3	7	6.4	59	15.0
Mesothelioma	0	0.0	0	0.0	0	0.0	0	0.1	1	0.0	0	0.0	0	0.0
Kaposi sarcoma	18	12.0	20	5.5	19	8.8	18	7.7	89	5.6	9	8.0	137	10.9
Connective and soft tissue	5	3.3	1	0.3	4	1.7	25	1.6	0	0.0	11	0.9	10	3.5
Breast	19	12.7	130	35.8	80	37.2	76	32.6	479	30.2	24	21.4	313	24.9
Vulva	2	1.3	4	1.1	1	0.5	2	0.9	15	0.9	0	0.0	2	1.3
Vagina	1	0.7	1	0.3	5	2.3	1	0.4	8	0.5	0	0.0	6	0.5
Cervix uteri	58	38.7	36	9.9	32	14.9	46	19.7	221	13.9	27	24.1	289	23.0
Corpus uteri	0	0.0	8	2.2	8	3.4	28	1.8	2	1.8	29	2.3	44	18.7
Uterus unspecified	2	1.3	2	0.6	2	0.9	0	0.0	9	0.6	0	0.0	7	0.7
Ovary	4	2.7	15											

Table 7: NAMIBIA (2010-2014) Basis of diagnosis of cancers – Males

SITE	Death certificate only		Clinical only		Clinical investigation		Biological/Immunological test			Haematology			Cytology			Histology of primary		Histology of secondary		Total				
	n	%	n	%	n	%	n	%	n	%	n	%	n	%	n	%	n	%	n	%	n	%		
Lip	0	0.0	0	0.0	0	0	0	0.0	0	0.0	0	0.0	0	0.0	6	100.0	0	0.0	6	100				
Tongue	0	0.0	0	0.0	0	0	0	0.0	0	0.0	0	0.0	1	1.0	102	99.0	0	0.0	103	100				
Mouth	1	0.9	0	0.0	0	0	0	0.0	0	0.0	2	1.7	113	97.4	0	0.0	116	100						
Salivary glands	1	2.4	0	0.0	0	0	0	0.0	1	2.4	0	0.0	0	0.0	39	95.1	0	0.0	41	100				
Tonsil	0	0.0	0	0.0	0	0	0	0.0	0	0.0	0	0.0	0	0.0	24	100.0	0	0.0	24	100				
Other oropharynx	0	0.0	0	0.0	0	0	0	0.0	0	0.0	0	0.0	0	0.0	11	100.0	0	0.0	11	100				
Nasopharynx	0	0.0	0	0.0	0	0	0	0.0	0	0.0	0	0.0	0	0.0	23	100.0	0	0.0	23	100				
Hypopharynx	0	0.0	0	0.0	0	0	0	0.0	0	0.0	0	0.0	0	0.0	6	100.0	0	0.0	6	100				
Pharynx unspecified	0	0.0	0	0.0	0	0	0	0.0	0	0.0	0	0.0	0	0.0	5	100.0	0	0.0	5	100				
Oesophagus	4	3.7	4	3.7	0	0	0	0.0	0	0.0	0	0.0	0	0.0	100	92.6	0	0.0	108	100				
Stomach	0	0.0	0	0.0	0	0	0	0.0	0	0.0	0	0.0	0	0.0	72	100.0	0	0.0	72	100				
Small intestine	0	0.0	0	0.0	0	0	0	0.0	0	0.0	0	0.0	0	0.0	6	100.0	0	0.0	6	100				
Colon	1	0.8	1	0.8	0	0	0	0.0	1	0.8	1	0.8	125	96.9	0	0.0	129	100						
Rectum	0	0.0	1	1.4	0	0	0	0.0	0	0.0	0	0.0	0	0.0	68	98.6	0	0.0	69	100				
Anus	0	0.0	0	0.0	0	0	0	0.0	0	0.0	0	0.0	0	0.0	23	100.0	0	0.0	23	100				
Liver	1	0.8	16	12.9	0	0	16	12.9	1	0.8	0	0.0	0	0.0	90	72.6	0	0.0	124	100				
Gallbladder etc.	0	0.0	0	0.0	0	0	0	0.0	0	0.0	0	0.0	0	0.0	12	100.0	0	0.0	12	100				
Pancreas	2	7.1	5	17.9	0	0	4	14.3	0	0.0	1	3.6	16	57.1	0	0.0	28	100						
Nose, sinuses etc.	0	0.0	0	0.0	0	0	0	0.0	0	0.0	0	0.0	0	0.0	20	100.0	0	0.0	20	100				
Larynx	0	0.0	0	0.0	0	0	0	0.0	0	0.0	0	0.0	0	0.0	147	100.0	0	0.0	147	100				
Trachea, bronchus and lung	0	0.0	34	21.1	0	0	0	0.0	6	3.7	1	0.6	120	74.5	0	0.0	151	100						
Other thoracic organs	0	0.0	0	0.0	0	0	0	0.0	0	0.0	0	0.0	0	0.0	8	100.0	0	0.0	8	100				
Bone	0	0.0	0	0.0	0	0	0	0.0	0	0.0	0	0.0	0	0.0	72	100.0	0	0.0	72	100				
Melanoma of skin	0	0.0	0	0.0	0	0	0	0.0	0	0.0	1	1.1	87	98.9	0	0.0	88	100						
Other skin	0	0.0	0	0.0	0	0	0	0.0	0	0.0	0	0.0	0	0.0	337	99.7	1	0.3	338	100				
Mesothelioma	0	0.0	0	0.0	0	0	0	0.0	0	0.0	0	0.0	0	0.0	7	100.0	0	0.0	7	100				
Kaposi sarcoma	142	16.1	0	0.0	0	0	0	0.0	0	0.0	0	0.0	0	0.0	740	83.9	0	0.0	882	100				
Connective and soft tissue	0	0.0	2	2.2	0	0	0	0.0	0	0.0	0	0.0	0	0.0	89	97.8	0	0.0	91	100				
Breast	1	1.6	0	0.0	0	0	0	0.0	0	0.0	0	0.0	0	0.0	61	98.4	0	0.0	62	100				
Penis	0	0.0	0	0.0	0	0	0	0.0	0	0.0	0	0.0	0	0.0	52	100.0	0	0.0	52	100				
Prostate	1	0.1	8	0.7	0	0	43	3.8	2	0.2	1	0.1	1073	95.1	0	0.0	1128	100						
Testis	1	3.1	0	0.0	0	0	0	0.0	0	0.0	0	0.0	0	0.0	31	96.9	0	0.0	32	100				
Other male genital organs	0	0.0	0	0.0	0	0	0	0.0	0	0.0	0	0.0	0	0.0	10	100.0	0	0.0	10	100				
Kidney	0	0.0	4	4.5	0	0	0	0.0	0	0.0	0	0.0	0	0.0	84	95.5	0	0.0	88	100				
Renal pelvis	0	0.0	0	0.0	0	0	0	0.0	0	0.0	0	0.0	0	0.0	47	95.5	0	0.0	49	100				
Ureter	0	0.0	0	0.0	0	0	0	0.0	0	0.0	0	0.0	0	0.0	61	100.0	0	0.0	61	100				
Bladder	0	0.0	0	0.0	0	0	0	0.0	0	0.0	0	0.0	0	0.0	7	100.0	0	0.0	7	100				
Other urinary organs	0	0.0	0	0.0	0	0	0	0.0	0	0.0	0	0.0	0	0.0	72	100.0	0	0.0	72	100				
Eye	0	0.0	0	0.0	0	0	0	0.0	0	0.0	0	0.0	0	0.0	241	100.0	0	0.0	241	100				
Brain, nervous system	1	1.4	22	31.9	0	0	0	0.0	0	0.0	0	0.0	0	0.0	46	66.7	0	0.0	69	100				
Thyroid	0	0.0	0	0.0	0	0	0	0.0	0	0.0	0	0.0	0	0.0	21	100.0	0	0.0	21	100				
Adrenal gland	0	0.0	0	0.0	0	0	0	0.0	0	0.0	0	0.0	0	0.0	3	100.0	0	0.0	3	100				
Other endocrine	0	0.0	2	40.0	0	0	0	0.0	0	0.0	0	0.0	0	0.0	3	60.0	0	0.0	5	100				
Hodgkin disease	0	0.0	0	0.0	0	0	0	0.0	0	0.0	0	0.0	0	0.0	32	100.0	0	0.0	32	100				
Non-Hodgkin lymphoma	0	0.0	0	0.0	0	0	0	0.0	0	0.0	0	0.0	0	0.0	230	99.6	0	0.0	231	100				
Immunoproliferative diseases	0	0.0	0	0.0	0	0	0	0.0	0	0.0	0	0.0	0	0.0	2	100.0	0	0.0	2	100				
Multiple myeloma	0	0.0	4	16.7	0	0	2	8.3	1	4.3	0	0.0	0	0.0	17	79.8	0	0.0	24	100				
Lymphoid leukaemia	0	0.0	0	0.0	0	0	0	0.0	0	0.0	0	0.0	0	0.0	49	95.1	0	0.0	51	100				
Myeloid leukaemia	0	0.0	0	0.0	0	0	0	0.0	0	0.0	0	0.0	0	0.0	52	100.0	0	0.0	52	100				
Leukaemia unspecified	0	0.0	0	0.0	0	0	0	0.0	0	0.0	0	0.0	0	0.0	4	100.0	0	0.0	4	100				
Myeloproliferative disorders	0	0.0	0	0.0	0	0	0	0.0	0	0.0	0	0.0	0	0.0	4	80.0	0	0.0	5	100				
Myelodysplastic syndromes	0	0.0	0	0.0	0	0	0	0.0	0	0.0	0	0.0	0	0.0	5	100.0	0	0.0	5	100				
Other and unspecified	1	0.5	2	0.9	0	0	0	0.0	0	0.0	0	0.0	0	0.0	15	7.0	195	91.5	0	0.0	213	100		
All sites	157	3.1	105	2.0	0	0	65	1.3	15	0.3	24	0.5	4758	92.8	1	0.0	5125	100						
All sites but C44	157	3.3	105	2.2	0	0	65	1.4	15	0.3	24	0.5	4421	92.4	0	0.0	4787	100						

*all row %

Table 8: NAMIBIA (2010-2014) Basis of diagnosis of cancers – Females

SITE	Death certificate only		Clinical only		Clinical investigation		Biological/Immunological test			Haematology			Cytology			Histology of primary		Histology of secondary		Total		
	n	%	n	%	n	%	n	%	n	%	n	%	n	%	n	%	n	%	n	%	n	%
Lip	0	0.0	0	0.0	0	0	0	0.0	0	0.0	0	0.0	0	0.0	6	100.0	0	0.0	6	100		
Tongue	0	0.0	0	0.0	0	0	0	0.0	0	0.0	0	0.0	0	0.0	46	100.0	0	0.0	46	100		
Mouth	3	3.4	0	0.0	0	0	0	0.0	0	0.0</												

Appendix: NAMBIA Annual incidence per 100 000 by age group and sex for single years

Central Cancer Registry, NAMIBIA (2010)

2010 Population Namibia

Incidence per 100,000 by age group (Period) - Male

SITE	ALL AGE AGES UNK	Incidence per 100,000 by age group (Period) - Male														CRUDE RATE (%)	CUM CUM ASR ICD (10th)					
		0-	5-	10-	15-	20-	25-	30-	35-	40-	45-	50-	55-	60-	65-	70-	75-	80+				
Lip	0 0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.0 0.0 0.00 0.00 0.0 C00				
Tongue	19 0	-	-	-	-	-	-	-	-	10.4	24.6	4.6	10.8	14.8	9.9	14.5	8.7	1.8 1.9 0.25 0.38 3.3 C01-02				
Mouth	33 0	-	-	-	-	-	-	-	-	7.8	14.1	36.9	16.3	22.2	9.9	58.1	34.7	3.2 3.4 0.40 0.56 5.3 C03-06				
Salivary glands	5 0	-	-	-	-	-	-	-	-	-	3.5	4.6	10.8	-	-	-	-	0.5 0.5 0.09 0.09 0.9 C07-08				
Tonsil	4 0	-	-	-	-	-	-	-	-	-	7.0	-	5.4	-	-	-	-	0.4 0.4 0.06 0.11 0.8 C09				
Other oropharynx	4 0	-	-	-	-	-	-	-	-	-	-	9.2	-	-	-	-	-	0.4 0.4 0.05 0.05 0.7 C10				
Nasopharynx	3 0	-	-	-	-	-	-	-	-	-	5.2	-	-	-	-	-	-	0.3 0.3 0.03 0.06 0.5 C11				
Hypopharynx	1 0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.1 0.1 0.00 0.00 0.1 C12-13				
Pharynx unspecified	1 0	-	-	-	-	-	-	-	-	-	-	-	5.4	-	-	-	-	0.1 0.1 0.03 0.03 0.2 C14				
Oesophagus	24 0	-	0.8	-	-	-	-	1.3	-	-	2.6	7.0	9.2	27.1	-	88.7	-	2.3 2.4 0.24 0.68 4.2 C15				
Stomach	15 0	-	-	-	-	-	-	-	-	-	5.2	10.5	4.6	-	22.2	9.9	72.7	-	1.4 1.5 0.10 0.26 2.6 C16			
Small intestine	2 0	-	-	-	-	-	-	-	-	-	2.6	-	-	-	-	-	-	0.2 0.2 0.01 0.05 0.4 C17				
Colon	29 0	-	-	-	-	-	-	-	3.2	6.2	10.4	17.6	13.8	5.4	29.7	39.4	43.6	-	2.8 2.9 0.28 0.63 5.0 C18			
Rectum	13 0	-	-	-	-	-	-	-	-	-	2.1	7.8	-	4.6	5.4	14.8	19.7	29.1	8.7	1.2 1.3 0.10 0.27 2.2 C19-20		
Anus	3 0	-	-	-	-	-	-	-	-	-	2.1	2.6	3.5	-	-	-	-	0.3 0.3 0.04 0.04 0.5 C21				
Liver	30 0	-	-	0.8	-	0.9	2.2	-	-	8.3	5.2	17.6	23.1	16.3	37.1	9.9	14.5	-	2.9 3.0 0.37 0.61 5.0 C22			
Gallbladder etc.	1 0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.1 0.1 0.00 0.05 0.2 C23-24				
Pancreas	11 0	-	-	-	-	-	-	-	1.3	-	-	7.0	-	21.7	22.2	9.9	-	-	1.1 1.1 0.15 0.31 2.2 C25			
Nose, sinuses etc.	3 0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.3 0.3 0.05 0.05 0.6 C30-31				
Larynx	43 0	-	-	-	-	-	-	-	2.2	-	-	5.2	17.6	18.5	21.7	126.1	9.9	87.2	17.3	4.1 4.4 0.33 1.01 8.0 C32		
Trachea, bronchus and lung	44 0	-	-	-	-	-	-	-	-	4.8	8.3	10.4	7.0	46.2	43.4	29.7	59.1	14.5	17.3	4.2 4.5 0.60 1.04 7.7 C33-34		
Other thoracic organs	1 0	-	-	-	-	-	-	-	-	1.1	-	-	-	-	-	-	-	0.1 0.1 0.01 0.01 0.1 C37-38				
Bone	13 0	-	-	-	-	-	-	-	-	-	5.2	-	4.6	10.8	-	-	-	-	1.2 1.3 0.13 0.13 1.6 C40-41			
Melanoma of skin	17 0	-	-	-	-	-	-	-	4.0	1.6	-	2.6	14.1	9.2	-	-	-	-	1.6 1.7 0.16 0.16 2.3 C43			
Other skin	93 0	-	-	-	-	-	-	1.7	0.9	1.1	13.4	17.7	12.4	36.5	24.6	23.1	54.2	51.9	39.4	116.3	60.7	8.9 9.5 0.93 1.38 13.5 C44
Mesothelioma	0 0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.0 0.0 0.00 0.00 0.0 C45				
Kaposi sarcoma	183 0	-	1.3	0.8	-	-	-	2.8	11.2	53.7	49.7	66.2	44.3	70.3	23.1	37.9	44.5	39.4	72.7	-	17.5 18.6 1.81 2.23 23.0 C46	
Connective and soft tissue	10 0	-	-	-	-	-	-	0.8	-	1.3	-	-	2.6	3.5	4.6	16.3	7.4	-	14.5	-	1.0 1.0 0.15 0.18 1.7 C47,C49	
Breast	17 0	-	-	-	-	-	-	-	-	2.2	2.7	-	7.8	17.6	13.8	-	7.4	9.9	-	-	1.6 1.7 0.22 0.31 2.7 C50	
Penis	14 0	-	-	-	-	-	-	-	-	-	2.7	-	13.0	-	18.5	-	-	-	-	1.3 1.4 0.17 0.17 2.0 C60		
Prostate	223 0	-	-	-	-	-	-	-	-	1.3	-	4.1	2.6	14.1	50.8	195.2	304.1	443.4	508.7	407.4	21.3 22.7 1.34 5.08 38.2 C61	
Testis	9 0	-	-	0.8	-	-	-	-	-	1.1	5.4	1.6	-	5.2	-	-	-	-	0.9 0.9 0.07 0.07 0.9 C62			
Other male genital organs	1 0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.1 0.1 0.02 0.02 0.2 C63				
Kidney	12 0	-	-	0.8	0.8	-	-	-	-	-	1.6	-	-	-	-	-	-	-	1.1 1.2 0.14 0.17 1.9 C64			
Renal pelvis	1 0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.1 0.1 0.02 0.02 0.2 C65				
Uterus	0 0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.0 0.0 0.00 0.00 0.0 C66				
Bladder	11 0	-	0.7	-	-	-	-	-	-	-	2.1	-	-	-	-	-	-	-	1.1 1.1 0.09 0.18 1.7 C67			
Other urinary organs	0 0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.0 0.0 0.00 0.00 0.0 C68				
Eye	62 0	-	2.0	0.8	-	-	0.9	1.1	13.4	14.4	20.7	13.5	23.5	17.6	23.1	-	14.8	29.6	43.6	-	5.9 6.3 0.59 0.81 8.1 C69	
Brain, nervous system	20 0	-	-	-	-	-	-	3.3	1.7	3.7	1.1	-	-	5.2	7.0	4.6	10.8	-	19.7	-	1.9 2.0 0.19 0.29 2.5 C70-72	
Thyroid	8 0	-	-	-	-	-	-	-	-	-	2.1	5.2	-	18.5	-	-	-	-	0.8 0.8 0.13 0.13 1.3 C73			
Adrenal gland	1 0	-	-	0.8	-	-	-	-	-	-	2.6	-	-	-	-	-	-	-	0.1 0.1 0.00 0.00 0.1 C74			
Other endocrine	0 0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.0 0.0 0.00 0.00 0.0 C75				
Hodgkin disease	2 0	-	-	-	-	-	-	-	-	1.6	-	-	-	-	-	-	-	-	0.2 0.2 0.01 0.01 0.3 C81			
Non-Hodgkin lymphoma	35 0	-	0.7	-	0.8	-	-	2.2	2.7	3.2	4.1	28.7	7.0	13.8	21.7	29.7	9.9	-	-	3.3 3.6 0.42 0.62 5.5 C82-85,C96		
Immunoproliferative diseases	0 0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.0 0.0 0.00 0.00 0.0 C98				
Multiple myeloma	3 0	-	-	-	-	-	-	-	-	-	1.6	-	-	3.5	-	-	-	-	0.3 0.3 0.03 0.06 0.5 C90			
Lymphoid leukaemia	13 0	-	1.3	1.6	-	-	0.9	1.1	-	-	2.1	5.2	3.5	-	5.4	-	-	-	1.2 1.3 0.11 0.20 1.7 C91			
Myeloid leukaemia	6 0	-	-	-	-	-	-	0.9	2.2	-	-	-	-	3.5	-	-	-	-	0.6 0.6 0.03 0.13 0.8 C92-94			
Leukaemia unspecified	0 0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.0 0.0 0.00 0.00 0.0 C95				
Myeloproliferative disorders	0 0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.0 0.0 0.00 0.00 0.0 MPD				
Myelodysplastic syndromes	0 0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.0 0.0 0.00 0.00 0.0 MDS				
Other and unspecified	34 0	-	0.7	-	0.8	-	-	1.3	1.6	2.1	-	7.8	10.5	13.8	21.7	66.7	29.6	43.6	8.7	3.2 3.5 0.30 0.78 6.0 O&U		
All sites	1077 0	-	6.6	6.5	6.6	5.9	5.8	13.8	29.1	107.4	104.3	142.6	273.6	344.3	410.8	596.3	889.9	955.8	1322.7	684.8	102.9 102.4 19.47 167.1 ALL	
All sites but C44	984 0	-	6.6	6.5	6.6	4.2	12.9	28.0	94.0	86.7	130.2	237.1	319.7	387.7	542.1	838.0	916.3	1206.4	624.1	94.0 100.0 9.31 18.08 153.5 ALLbC44		

, Reference population: World Standard Population

Table builtFri Nov 18 07:49:39 SAST 2016 by CanReg5.

Central Cancer Registry, NAMIBIA (2010)

2010 Population Namibia

Incidence per 100,000 by age group (Period) - Female

SITE	ALL AGE AGES UNK	Incidence per 100,000 by age group (Period) - Female														CRUDE RATE (%)	CUM CUM ASR ICD (10th)		
		0-	5-	10-	15-	20-	25-	30-	35-	40-	45-	50-	55-	60-	65-	70-	75-	80+	
Lip	0 0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.0 0.0 0.00 0.00 0.0 C00	
Tongue	12 0	-	-	-	-	-	-	-	-	3.8	-	11.3	-	4.4	7.2	9.9	4.6	-	1.1 1.1 0.10 0.16 1.5 C01-02
Mouth	25 0	-	-																

Central Cancer Registry, NAMIBIA (2011)

2011 Population Namibia

Incidence per 100,000 by age group (Period) - Male

Site	All age																					Crude Rate	(%)	Cum 0-64	Cum 0-74	ASR	ICD (10th)
	Age unk	0-	5-	10-	15-	20-	25-	30-	35-	40-	45-	50-	55-	60-	65-	70-	75-	80+									
Lip	1	0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	8.7	-	0.1	0.1	0.00	0.00	0.1	C00	
Tongue	17	0	-	-	-	-	-	-	-	2.1	-	21.1	9.2	10.8	14.8	19.7	14.5	-	8.7	-	1.6	2.0	0.22	0.39	3.1	C01-02	
Mouth	14	0	-	-	-	-	-	-	-	-	7.8	7.0	4.6	5.4	22.2	-	-	-	17.3	-	1.3	1.6	0.14	0.25	2.2	C03-06	
Salivary glands	9	0	-	-	-	-	-	-	-	-	-	13.0	7.0	-	-	-	-	-	-	29.1	-	0.9	1.1	0.10	0.10	1.4	C07-08
Tonsil	2	0	-	-	-	-	-	-	-	-	-	2.6	-	-	-	-	-	-	9.9	-	-	0.2	0.2	0.01	0.06	0.4	C09
Other oropharynx	0	0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.0	0.0	0.00	0.00	0.0	C10	
Nasopharynx	4	0	-	0.8	-	-	-	0.9	1.1	-	-	2.1	-	-	-	-	-	-	-	-	0.4	0.5	0.02	0.02	0.4	C11	
Hypopharynx	2	0	-	-	-	-	-	-	-	-	-	-	-	-	4.6	5.4	-	-	-	-	0.2	0.2	0.05	0.05	0.4	C12-13	
Pharynx unspecified	1	0	-	-	-	-	-	-	1.3	-	-	-	-	-	-	-	-	-	14.5	-	0.1	0.1	0.01	0.01	0.1	C14	
Oesophagus	19	0	-	-	-	-	-	-	-	1.6	-	2.6	10.5	13.8	16.3	37.1	19.7	14.5	-	-	1.8	2.2	0.22	0.51	3.6	C15	
Stomach	12	0	-	-	-	-	-	-	-	6.2	-	-	-	-	-	29.7	9.9	43.6	8.7	-	1.1	1.4	0.03	0.23	2.0	C16	
Small intestine	1	0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	14.5	-	0.1	0.1	0.00	0.00	0.1	C17	
Colon	17	0	-	-	-	-	-	0.9	1.1	4.0	1.6	2.1	7.8	10.5	-	-	-	-	29.7	-	1.6	2.0	0.14	0.29	2.5	C18	
Rectum	12	0	-	-	-	-	-	-	-	-	-	6.4	-	3.5	4.6	5.4	7.4	-	-	43.6	8.7	1.1	1.4	0.07	0.14	2.0	C19-20
Anus	5	0	-	-	-	-	-	-	-	1.1	-	-	5.2	7.0	-	-	-	-	-	-	0.5	0.6	0.07	0.07	0.8	C21	
Liver	28	0	-	0.7	-	-	-	-	-	5.4	4.8	6.2	2.6	7.0	27.7	10.8	14.8	19.7	14.5	8.7	2.7	3.3	0.33	0.50	4.2	C22	
Gallbladder etc.	1	0	-	-	-	-	-	-	-	-	-	-	-	3.5	4.6	-	-	-	9.9	14.5	-	0.1	0.1	0.02	0.02	0.2	C23-24
Pancreas	4	0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.4	0.5	0.04	0.09	0.7	C25	
Nose, sinuses etc.	4	0	-	-	-	-	-	-	-	1.6	-	2.6	7.0	-	-	-	-	-	-	-	0.4	0.5	0.06	0.06	0.6	C30-31	
Larynx	26	0	-	-	-	-	-	1.1	-	-	-	7.8	17.6	18.5	10.8	37.1	19.7	29.1	17.3	-	2.5	3.0	0.28	0.56	4.6	C32	
Trachea, bronchus and lung	17	0	-	-	-	-	-	-	-	3.2	-	-	3.5	23.1	10.8	29.7	9.9	14.5	8.7	-	1.6	2.0	0.20	0.40	3.0	C33-34	
Other thoracic organs	3	0	1.3	-	-	-	-	-	-	-	-	-	-	4.6	-	-	-	-	-	-	0.3	0.4	0.03	0.03	0.3	C37-38	
Bone	8	0	-	1.6	-	1.7	-	1.1	1.3	1.6	-	-	-	-	-	-	-	9.9	-	-	0.8	0.9	0.04	0.09	0.8	C40-41	
Melanoma of skin	20	0	-	-	-	-	-	-	1.1	-	3.2	-	2.6	3.5	13.8	10.8	44.5	-	14.5	26.0	1.9	2.3	0.18	0.40	3.3	C43	
Other skin	37	0	-	-	-	-	-	6.7	2.7	8.0	4.1	10.4	10.5	13.8	-	7.4	29.6	58.1	34.7	-	3.5	4.3	0.28	0.47	4.9	C44	
Mesothelioma	1	0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.1	0.1	0.00	0.00	0.1	C45	
Kaposi sarcoma	200	0	-	-	-	2.5	3.4	1.8	12.3	61.8	46.5	66.2	78.2	49.2	50.8	43.4	22.2	29.6	29.1	17.3	19.1	23.4	2.08	2.34	2.48	C46	
Connective and soft tissue	18	0	-	1.3	-	-	-	3.7	-	2.7	1.6	-	-	3.5	13.8	5.4	14.8	-	14.5	8.7	1.7	2.1	0.16	0.23	2.3	C47-C49	
Breast	16	0	-	-	-	-	-	-	-	4.0	1.6	4.1	2.6	3.5	9.2	5.4	7.4	-	43.6	8.7	1.5	1.9	0.15	0.19	2.2	C50	
Penis	7	0	-	-	-	-	-	-	1.1	1.3	1.6	2.1	-	-	-	-	-	14.8	-	14.5	-	0.7	0.8	0.03	0.10	1.0	C60
Prostate	221	1	-	-	-	-	-	-	1.3	1.6	2.1	7.8	14.1	32.3	146.4	318.9	463.1	494.2	450.7	-	21.1	25.8	1.03	4.90	37.1	C61	
Testis	2	0	-	-	-	-	-	-	1.1	-	-	-	-	-	-	-	-	-	-	-	0.2	0.2	0.01	0.01	0.2	C62	
Other male genital organs	2	0	-	-	-	-	-	-	0.9	-	-	-	-	-	-	-	-	-	-	-	0.2	0.2	0.00	0.00	0.2	C63	
Kidney	15	0	2.6	1.6	-	-	-	-	-	2.1	5.2	-	13.8	-	7.4	-	14.5	8.7	-	1.4	1.8	0.18	0.36	1.9	C64		
Urinary pelvis	0	0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.0	0.0	0.00	0.00	0.0	C65	
Uterus	0	0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.0	0.0	0.00	0.00	0.0	C66	
Bladder	9	0	-	-	-	-	-	-	-	-	-	-	3.5	4.6	5.4	37.1	9.9	-	-	14.5	-	0.9	1.1	0.07	0.30	1.9	C67
Other urinary organs	1	0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	14.5	-	0.1	0.1	0.00	0.00	0.1	C68	
Eye	45	0	4.0	-	-	-	2.2	6.7	12.8	12.4	15.6	3.5	13.8	10.8	29.7	9.9	-	-	8.7	-	4.3	5.3	0.41	0.61	5.8	C69	
Brain, nervous system	13	0	2.6	-	0.8	-	-	-	2.7	-	2.7	2.6	3.5	9.2	-	-	-	-	-	-	1.2	1.5	0.12	0.12	1.5	C70-72	
Thyroid	3	0	-	-	-	-	-	-	-	3.2	-	-	-	-	-	-	-	-	-	-	0.3	0.4	0.03	0.03	0.3	C73	
Adrenal gland	1	0	-	0.7	-	-	-	-	-	-	2.1	-	-	-	-	-	-	-	-	-	0.1	0.1	0.00	0.00	0.1	C74	
Other endocrine	2	0	-	-	-	-	-	-	-	-	1.6	-	-	-	5.4	-	-	-	-	-	0.2	0.2	0.04	0.04	0.3	C75	
Hodgkin disease	7	0	0.7	0.8	0.8	-	1.1	-	3.2	-	-	-	4.6	-	-	-	-	-	-	0.7	0.8	0.06	0.06	0.7	C81		
Non-Hodgkin lymphoma	26	0	0.7	-	0.8	0.8	-	1.3	3.2	12.4	7.8	3.5	18.5	5.4	7.4	-	-	-	8.7	-	2.5	3.0	0.29	0.33	3.4	C82-85,C96	
Immunoproliferative diseases	0	0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.0	0.0	0.00	0.00	0.0	C88	
Lymphoid leukaemia	6	0	-	0.8	-	0.8	-	-	-	-	-	-	4.6	5.4	-	9.9	14.5	-	-	0.6	0.7	0.06	0.11	0.9	C91		
Myeloid leukaemia	5	0	0.7	-	0.8	-	-	-	-	-	-	5.2	-	-	7.4	-	-	-	-	-	0.5	0.6	0.03	0.07	0.7	C92-94	
Leukaemia unspecified	0	0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.0	0.0	0.00	0.00	0.0	C95	
Myeloproliferative disorders	1	0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.1	0.1	0.00	0.05	0.2	MPD	
Melodysplastic syndromes	0	0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.0	0.0	0.00	0.00	0.0	MDS	
Other and unspecified	23	0	-	-	-	-	-	-	1.1	4.0	3.2	2.1	5.2	-	13.8	16.3	22.2	29.6	14.5	8.7	2.2	2.7	0.23	0.49	3.7	O&U	
All sites	893	1	15.2	4.8	5.8	7.6	9.2	35.8	102.0	112.3	130.2	195.4	207.3	341.6	341.5	787.8	719.3	973.8	693.4	-	85.3	7.55	15.05	131.8	ALL		
All sites but C44	856	1	15.2	4.8	5.8	7.6	9.2	29.1	99.4	104.3	126.1	185.0	196.8	327.7	341.5	771.3	689.7	915.7	658.8	-	81.8	10.00	7.27	14.58	126.9	ALLbC44	

, Reference population: World Standard Population

Table built Fri Nov 18 07:50:19 SAST 2016 by CanReg5.

Central Cancer Registry, NAMIBIA (2011)

2011 Population Namibia

Incidence per 100,000 by age group (Period) - Female

Site	All Age Ages Unk													Crude Rate (%)	Cum (%) 0-64	Cum (%) 65-74	ASR ICD (10th)			
	0-	5-	10-	15-	20-	25-	30-	35-	40-	45-	50-	55-	60-	65-	70-	75-	80+			
Lip	0	0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.0 0.0 0.00 0.0 C00		
Tongue	6	0	0.7	-	-	-	-	-	-	2.3	2.8	-	-	-	-	-	-	0.5 0.6 0.03 0.06 C01-02		
Mouth	16	0	-	-	-	-	-	-	-	2.3	8.5	11.0	4.4	27.3	7.2	-	-	1.4 1.7 0.13 0.30 C03-06		
Salivary glands	3	0	-	-	-	-	-	-	-	2.3	2.8	-	-	-	-	-	-	0.3 0.3 0.03 0.03 C07-08		
Tonsil	2	0	-	-	-	-	-	-	-	-	3.7	-	-	-	-	-	-	0.2 0.2 0.02 0.02 C09		
Other oropharynx	0	0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.0 0.0 0.00 0.00 C10		
Nasopharynx	3	0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.3 0.3 0.02 0.08 C11		
Hypopharynx	0	0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.0 0.0 0.00 0.00 C12-13		
Pharynx unspecified	0	0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.0 0.0 0.00 0.00 C14		
Oesophagus	7	0	-	-	-	-	-	1.3	-	1.9	2.3	-	7.4	4.4	-	-	4.6	0.6 0.7 0.09 0.09 C15		
Stomach	6	0	-	-	-	-	-	-	-	2.3	-	7.4	8.8	-	-	-	4.6	0.5 0.6 0.09 0.09 C16		
Small intestine	0	0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.0 0.0 0.00 0.00 C17		
Colon	24	0	-	0.8	-	0.9	-	1.3	-	5.7	2.3	14.1	11.0	8.8	10.9	14.3	19.8	4.6	2.2 2.5 0.22 0.35 C18	
Rectum	14	0	-	-	-	-	-	1.1	2.6	3.0	1.9	6.8	-	8.8	5.5	-	9.9	4.6	1.3 1.7 0.15 0.25 C19-20	
Anus	3	0	-	-	-	-	-	1.1	1.3	1.5	-	-	-	-	-	-	-	0.3 0.3 0.02 0.02 C21		
Liver	11	0	-	-	-	-	-	0.9	-	1.3	1.5	-	2.8	-	17.7	5.5	7.2	-	1.0 1.2 0.12 0.18 C22	
Gallbladder etc.	2	0	-	-	-	-	-	-	-	-	-	2.3	2.8	-	-	-	-	0.2 0.2 0.03 0.03 C23-24		
Pancreas	1	0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.1 0.1 0.00 0.04 C25		
Nose, sinuses etc.	3	0	-	-	-	-	-	-	2.6	-	-	-	-	-	-	-	-	0.3 0.3 0.01 0.04 C30-31		
Larynx	6	0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.5 0.6 0.02 0.11 C32		
Trachea, bronchus and lung	15	0	-	-	-	-	-	0.9	-	-	1.9	6.8	2.8	3.7	17.7	10.9	-	9.9	4.6	1.4 1.6 0.17 0.22 C33-34
Other thoracic organs	0	0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.0 0.0 0.00 0.00 C37-38		
Bone	14	0	-	0.8	3.3	3.3	0.9	-	-	1.5	1.9	2.3	-	-	-	-	7.2	-	1.3 1.5 0.07 0.11 C40-41	
Melanoma of skin	24	0	-	-	-	-	-	-	1.3	1.5	-	9.1	5.7	3.7	8.8	21.8	14.3	9.9	27.7	2.2 2.5 0.15 0.33 C43
Other skin	63	0	-	-	-	4.1	2.7	3.2	5.2	3.0	15.2	9.1	14.1	14.7	13.3	27.8	14.3	89.1	32.3	5.7 6.7 0.42 0.60 C44
Mesothelioma	0	0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.0 0.0 0.00 0.00 C45		
Kaposi sarcoma	116	0	-	1.6	4.1	2.5	4.5	8.6	31.0	41.2	24.7	27.2	17.0	14.7	4.4	16.4	7.2	-	10.5 12.3 0.91 1.02 C46	
Connective and soft tissue	17	0	2.0	0.8	0.8	-	-	2.1	2.6	-	1.9	4.5	2.8	3.7	8.8	5.5	-	-	1.5 1.8 0.15 0.18 C47,C49	
Breast	265	0	-	-	-	0.9	8.6	22.0	30.5	64.6	63.0	62.2	114.1	119.3	152.8	150.4	69.3	96.8	23.9 28.0 2.43 3.94 33.3 C50	
Vulva	6	0	-	-	-	-	-	-	1.9	2.3	2.8	-	4.4	5.5	-	9.9	-	-	0.5 0.6 0.06 0.08 0.8 C51	
Vagina	10	0	-	-	-	-	-	-	1.5	3.8	2.3	-	3.7	4.4	-	14.3	-	9.2	0.9 1.1 0.08 0.15 1.2 C52	
Cervix uteri	156	0	-	-	-	0.9	2.1	16.8	24.4	36.1	36.3	59.4	47.8	48.6	49.1	93.1	79.2	64.5	14.1 16.5 1.36 2.07 18.7 C53	
Cuprus uteri	17	0	-	-	-	-	-	-	1.3	-	1.9	-	5.7	11.0	13.3	16.4	7.2	9.9	1.5 1.8 0.17 0.28 2.3 C54	
Uterus unspecified	7	0	-	-	-	-	-	-	1.3	1.5	-	4.5	-	7.4	-	7.2	-	-	0.6 0.7 0.07 0.11 0.9 C55	
Ovary	27	0	-	-	0.8	1.6	1.8	1.1	2.6	-	-	2.3	19.8	7.4	8.8	10.9	14.3	29.7	-	2.4 2.9 0.23 0.36 3.3 C56
Other female genital organs	1	0	-	-	-	-	-	-	-	-	-	-	2.8	-	-	-	-	-	0.1 0.1 0.01 0.01 0.1 C57	
Placenta	6	0	-	-	-	-	-	2.7	1.1	-	1.9	-	3.7	-	-	-	-	-	0.5 0.6 0.05 0.05 0.6 C58	
Kidney	9	0	2.0	1.6	-	0.8	-	-	1.3	-	-	7.4	-	-	-	-	-	-	0.8 1.0 0.07 0.07 0.9 C64	
Renal pelvis	0	0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.0 0.0 0.00 0.00 0.0 C65		
Ureter	0	0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.0 0.0 0.00 0.00 0.0 C66		
Bladder	8	0	-	-	-	-	-	-	1.1	-	-	1.9	-	-	-	5.5	7.2	29.7	4.6	0.7 0.8 0.01 0.08 0.8 C67
Other urinary organs	1	0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.1 0.1 0.00 0.00 0.1 C68		
Eye	45	0	4.7	-	-	-	6.4	10.3	13.7	7.6	9.1	5.7	-	-	-	14.3	9.9	9.2	4.1 4.8 0.29 0.36 4.3 C69	
Brain, nervous system	11	0	-	0.8	1.6	-	0.9	2.1	-	-	2.3	2.8	7.4	4.4	-	-	-	-	1.0 1.2 0.11 0.17 1.2 C70,C72	
Thyroid	13	0	-	-	-	1.6	-	-	2.6	1.5	7.6	9.1	-	-	-	-	-	-	1.2 1.4 0.11 0.14 1.4 C73	
Adrenal gland	1	0	0.7	-	-	-	-	-	-	-	2.3	-	-	-	-	-	-	-	0.1 0.1 0.01 0.01 0.1 C74	
Other endocrine	1	0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.1 0.1 0.01 0.01 0.1 C75		
Hodgkin disease	3	0	-	-	-	-	0.9	1.1	-	-	1.9	-	-	-	-	-	-	-	0.3 0.3 0.02 0.02 0.3 C81	
Non-Hodgkin lymphoma	26	0	-	1.6	-	-	0.9	2.1	3.9	1.5	5.7	9.1	-	3.7	8.8	10.9	-	-	2.3 2.8 0.19 0.24 2.7 C82-85,C96	
Immunoproliferative diseases	0	0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.0 0.0 0.00 0.00 0.0 C88		
Multiple myeloma	10	0	-	-	-	-	-	-	1.3	-	-	-	-	14.7	4.4	10.9	-	9.9	4.6	0.9 1.1 0.10 0.16 1.3 C90
Lymphoid leukaemia	5	0	0.7	-	-	-	1.1	-	-	-	2.3	-	-	4.4	-	7.2	-	-	0.5 0.5 0.04 0.08 0.6 C91	
Myeloid leukaemia	6	0	-	-	0.8	-	0.9	1.1	-	-	4.5	-	3.7	-	-	-	-	-	0.5 0.6 0.06 0.06 0.7 C92-94	
Leukemia unspecified	0	0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.0 0.0 0.00 0.00 0.0 C95		
Myeloproliferative disorders	1	0	-	-	-	-	-	-	-	-	-	-	-	-	-	5.5	-	-	0.1 0.1 0.03 0.03 0.2 MPD	
Myelodysplastic syndromes	0	0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.0 0.0 0.00 0.00 0.0 MDS		
Other and unspecified	17	0	-	-	0.8	-	-	1.1	1.3	3.0	3.8	2.3	5.7	7.4	4.4	10.9	7.2	9.9	-	1.5 1.8 0.15 0.24 2.1 O&U
All sites	1008	0	10.8	7.4	13.1	13.9	20.6	45.0	114.9	131.1	193.7	233.8	242.3	320.2	340.2	436.6	408.3	405.9	354.9	90.9 8.44 12.66 117.6 ALL
All sites but C44	945	0	10.8	7.4	13.1	9.8	17.9	41.8	109.8	128.1	178.5	224.7	229.0	305.5	326.9	414.8	393.9	316.8	322.6	85.2 100.0 8.02 12.06 10.8 ALLbC44

, Reference population: World Standard Population

Table built Fri Nov 18 07:50:19 SAST 2016 by CanReg5.

Central Cancer Registry, NAMIBIA (2012)

2012 Population Namibia

Incidence per 100,000 by age group (Period) - Male

SITE	ALL AGE AGES UNK	0-	5-	10-	15-	20-	25-	30-	35-	40-	45-	50-	55-	60-	65-	70-	75-	80+	CRUDE RATE (%)	CUM CUM ASR ICD (0-64 0-74 (10th)					
Lip	0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.0	0.0 0.00 0.00 0.0 C00						
Tongue	13	0	-	-	-	-	-	-	-	-	2.6	7.0	13.8	10.8	14.8	19.7	14.5	-	1.2	1.5 0.17 0.34 2.5 C01-02					
Mouth	21	0	-	-	-	-	-	-	-	-	-	18.5	10.8	29.7	26.0	-	-	-	2.0	2.5 0.15 0.44 3.6 C03-06					
Salivary glands	9	0	-	-	-	-	-	2.2	1.3	1.6	-	2.6	-	13.8	-	7.4	-	-	0.9	1.1 0.11 0.15 1.3 C07-08					
Tonsil	8	0	-	-	-	-	-	-	2.2	-	-	2.6	3.5	-	5.4	-	-	29.1	0.8 0.04 0.04 1.1 C09						
Other oropharynx	3	0	-	-	-	-	-	-	-	1.6	-	-	-	5.4	-	-	14.5	0.3 0.4 0.04 0.04 0.5 C10							
Nasopharynx	7	0	-	-	-	-	-	0.8	-	1.1	-	-	-	5.2	3.5	-	-	-	0.7 0.8 0.05 0.13 1.1 C11						
Hypopharynx	1	0	-	-	-	-	-	-	-	-	-	-	-	3.5	-	-	-	-	0.1 0.1 0.02 0.02 0.2 C12-13						
Pharynx unspecified	0	0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.0 0.0 0.00 0.00 0.0 C14						
Oesophagus	17	0	-	-	-	-	-	-	-	1.3	-	2.1	5.2	7.0	18.5	16.3	14.8	9.9	14.5	1.6 2.0 0.25 0.38 3.0 C15					
Stomach	11	0	-	-	-	-	-	-	-	-	-	2.1	10.4	3.5	4.6	-	7.4	9.9	29.1	1.1 1.3 0.10 0.19 1.8 C16					
Small intestine	1	0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.1 0.1 0.02 0.02 0.2 C17						
Colon	19	0	-	-	-	-	-	-	-	1.1	-	-	2.1	2.6	17.6	13.8	16.3	14.8	-	1.8 2.2 0.27 0.34 3.3 C18					
Rectum	14	0	-	-	-	-	-	-	-	0.9	-	-	1.6	8.3	-	3.5	-	10.8	29.6	-	1.3 1.6 0.13 0.35 2.3 C19-20				
Anus	6	0	-	-	-	-	-	-	-	-	-	-	-	10.5	-	7.4	9.9	-	8.7	0.6 0.7 0.05 0.14 1.1 C21					
Liver	17	0	0.7	-	-	-	-	-	-	1.1	-	-	-	13.0	10.5	9.2	10.8	14.8	-	1.6 2.0 0.20 0.30 2.9 C22					
Gallbladder etc.	3	0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.3 0.4 0.04 0.00 0.3 C23-24						
Pancreas	5	0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.5 0.6 0.08 0.08 0.8 C25						
Nose, sinuses etc.	3	0	-	-	-	-	-	-	-	0.9	-	-	1.6	-	-	-	-	-	0.3 0.4 0.01 0.05 0.4 C30-31						
Larynx	26	0	-	-	-	-	-	-	-	-	-	-	-	4.1	-	3.5	13.8	27.1	22.2	59.1	43.6	26.0			
Trachea, bronchus and lung	28	0	-	-	-	-	-	-	-	-	-	-	-	2.1	2.6	7.0	23.1	21.7	59.3	39.4	29.1	8.7			
Other thoracic organs	0	0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.0 0.0 0.00 0.00 0.0 C37-38						
Bone	9	0	-	-	-	-	-	-	-	0.8	0.8	0.9	-	2.7	-	2.1	-	-	-	0.9 1.1 0.05 0.14 1.1 C40-41					
Melanoma of skin	14	0	-	-	-	-	-	-	-	-	-	-	-	2.7	-	7.0	9.2	10.8	-	1.3 1.6 0.15 0.20 1.9 C43					
Other skin	59	0	-	-	-	-	-	-	-	1.7	3.7	2.2	9.4	3.2	6.2	18.2	10.5	23.7	32.5	37.1	78.8	29.1	26.0		
Mesothelioma	1	0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.1 0.1 0.01 0.01 0.1 C45						
Kaposi's sarcoma	184	0	-	-	-	-	-	-	-	2.4	-	1.7	2.8	6.7	48.3	54.6	78.6	59.9	42.2	32.3	37.9	44.5	29.6	29.1	17.3
Connective and soft tissue	17	0	-	-	-	-	-	-	-	1.6	-	0.8	1.8	1.1	4.0	4.8	-	-	3.5	-	10.8	-	-	1.6 2.0 0.14 0.23 2.0 C47,C49	
Breast	9	0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.9 1.1 0.07 0.20 1.4 C50					
Penis	5	0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.5 0.6 0.04 0.04 0.7 C60						
Prostate	196	1	-	-	-	-	-	-	-	-	-	-	-	1.3	-	10.4	28.1	46.2	157.2	252.2	453.2	343.7	346.7	18.7 23.0 1.22 4.77 33.9 C61	
Testis	4	0	-	-	-	-	-	-	-	-	-	-	-	1.3	-	4.1	2.6	-	-	-	-	-	0.4 0.5 0.04 0.04 0.5 C62		
Other male genital organs	1	0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.1 0.1 0.02 0.02 0.2 C63					
Kidney	21	0	0.2	2.6	-	2.5	-	-	-	-	-	-	-	2.1	-	7.0	13.8	16.3	7.4	19.7	14.5	8.7	-	0.5 0.6 0.04 0.04 0.6 C64	
Renal pelvis	0	0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.0 0.0 0.00 0.00 0.0 C66					
Uterus	0	0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.0 0.0 0.00 0.00 0.0 C68					
Bladder	10	0	-	-	-	-	-	-	-	0.8	-	-	0.9	-	-	-	-	-	-	-	-	-	1.0 1.2 0.06 0.19 1.5 C67		
Other urinary organs	0	0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.0 0.0 0.00 0.00 0.0 C68					
Eye	48	0	2.6	-	-	-	-	-	-	3.4	-	8.1	9.6	16.5	26.1	3.5	4.6	10.8	22.2	-	-	-	4.6 5.6 0.44 0.55 6.0 C69		
Brain, nervous system	8	0	-	-	-	-	-	-	-	0.8	-	1.1	1.6	2.1	-	3.5	-	7.4	-	-	-	-	0.8 0.9 0.09 0.13 1.2 C70-72		
Thyroid	2	0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.2 0.2 0.02 0.06 0.4 C73					
Adrenal gland	1	0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.1 0.1 0.02 0.02 0.2 C74					
Other endocrine	1	0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.1 0.1 0.01 0.01 0.1 C75					
Hodgkin disease	7	0	-	-	-	-	-	-	-	0.8	1.8	-	1.3	1.6	4.1	-	-	-	-	-	-	0.7 0.8 0.05 0.05 0.6 C81			
Non-Hodgkin lymphoma	40	0	0.7	0.8	1.7	0.8	-	-	-	1.1	-	-	3.2	10.3	7.8	38.6	18.5	10.8	-	29.6	29.1	17.3	3.8 4.7 0.47 0.62 5.9 C82-85,C96		
Immunoproliferative diseases	0	0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.0 0.0 0.00 0.00 0.0 C88					
Multiple myeloma	4	0	-	-	-	-	-	-	-	1.8	-	-	-	-	-	-	-	-	-	-	-	0.4 0.5 0.04 0.09 0.6 C90			
Lymphoid leukaemia	9	0	1.3	0.8	0.8	-	-	-	-	-	-	-	-	2.1	-	7.0	-	-	-	-	-	-	0.9 1.1 0.06 0.06 1.0 C91		
Myeloid leukaemia	10	0	1.3	-	-	-	-	-	-	2.2	-	1.3	-	2.1	-	3.5	-	5.4	7.4	9.9	-	-	1.0 1.2 0.08 0.17 1.4 C92-94		
Leukaemia unspecified	1	0	-	-	-	-	-	-	-	0.8	-	-	-	-	-	-	-	-	-	-	-	0.1 0.1 0.00 0.00 0.1 C95			
Myeloproliferative disorders	0	0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.0 0.0 0.00 0.00 0.0 MPD					
Myelodysplastic syndromes	2	0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.2 0.2 0.02 0.06 0.4 MDS					
Other and unspecified	35	0	1.3	-	-	-	-	-	-	1.3	3.2	2.1	15.6	3.5	3.5	9.2	48.8	29.7	29.6	43.6	8.7	-	3.3 4.1 0.43 0.72 6.0 O&U		
All sites	910	1	10.6	6.5	6.6	12.7	15.7	25.7	85.9	96.3	155.0	187.6	256.5	323.1	504.1	689.7	926.2	799.4	632.7	-	-	-	87.0 84.4 16.53 13.81 ALL ALL		
All sites but C44	851	1	10.6	6.5	6.6	11.0	12.0	23.5	76.5	93.1	148.8	169.4	246.0	300.0	471.6	652.6	847.4	770.3	606.7	-	-	-	81.3 100.0 7.89 15.40 129.2 ALLbC44		

, Reference population: World Standard Population

Table builtFri Nov 18 07:55:48 SAST 2016 by CanReg5.

Central Cancer Registry, NAMIBIA (2012)

2012 Population Namibia

Incidence per 100,000 by age group (Period) - Female

SITE	ALL AGE AGES UNK	0-	5-	10-	15-	20-	25-	30-	35-	40-	45-	50-	55-	60-	65-	70-	75-	80+	CRUDE RATE (%)	CUM CUM ASR ICD (0-64 0-74 (10th)	
Lip	1	0	-	-	-	-	-	-	-	1.9	-	2.3	5.7	7.4	13.3	5.5	7.2	-	-	0.1 0.1 0.01 0.01 1.0 C00	
Tongue	11	0	-	-	-	-	-	-	-	1.9	-	2.3	5.7	18.4	8.8	10.9	-	9.9	-	-	1.0 1.5 0.15 0.22 1.7 C01-02
Mouth	14	0	-	-	-	-	-	-	-	1.5	-	2.3	5.7	18.4	8.8	10.9	-	9.9	-	-	1.3 1.5 0.18 0.24 2.0 C03-06
Salivary glands	4	0	-	-	-	-	-	-	-	-	-	-	-	-	5.5	-	-	-	-	0.4 0.4 0.01 0.04 0.4 C07-08	

Central Cancer Registry, NAMIBIA (2013)

2013 Population Namibia

Incidence per 100,000 by age group (Period) - Male

SITE	ALL AGES UNK	Incidence per 100,000 by age group (Period) - Male													CRUDE RATE (%)	CUM CUM ASR (0-64)	CUM CUM ASR (65-80+)	ICD (10th)				
		0-	5-	10-	15-	20-	25-	30-	35-	40-	45-	50-	55-	60-	65-	70-	75-	80+				
Lip	2 0	-	-	-	-	0.8	-	-	1.6	-	-	-	-	-	-	-	-	0.2	0.2 0.01 0.01	0.2 C00		
Tongue	23 0	-	-	-	-	0.8	-	0.9	1.1	-	-	4.1	2.6	14.1	4.6	27.1	7.4	39.4	14.5	26.0	2.2 0.2 0.27 0.50	3.9 C01-02
Mouth	24 0	-	-	-	-	0.8	-	1.1	-	-	-	2.6	14.1	-	10.8	37.1	39.4	14.5	34.7	2.3 2.3 0.15 0.53	3.9 C03-06	
Salivary glands	7 0	-	-	-	-	-	-	-	1.3	1.6	-	2.6	-	-	-	7.4	29.6	-	-	0.7 0.7 0.03 0.21	1.1 C07-08	
Tonsil	5 0	-	-	-	-	-	-	-	1.6	-	-	-	4.6	5.4	7.4	9.9	-	-	-	0.5 0.5 0.06 0.14	0.9 C09	
Other oropharynx	1 0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.1 0.1 0.00 0.05	0.2 C10		
Nasopharynx	3 0	-	-	-	-	-	-	-	-	-	-	3.5	4.6	-	-	-	-	8.7	-	0.3 0.3 0.04 0.04	0.4 C11	
Hypopharynx	0 0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.0 0.0 0.00 0.00	0.0 C12-13		
Pharynx unspecified	1 0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	9.9	-	-	-	0.1 0.1 0.00 0.05	0.2 C14	
Oesophagus	17 0	-	-	-	-	-	-	-	1.6	2.1	2.6	7.0	13.8	16.3	37.1	-	14.5	-	-	1.6 1.6 0.22 0.40	3.2 C15	
Stomach	18 0	-	-	-	-	-	-	-	1.6	2.1	2.6	10.5	4.6	5.4	7.4	29.6	43.6	26.0	-	1.7 1.7 0.13 0.32	2.8 C16	
Small intestine	1 0	-	-	-	-	-	-	-	-	-	-	5.4	-	-	-	-	-	-	0.1 0.1 0.03 0.03	0.2 C17		
Colon	37 0	-	-	-	-	-	-	1.1	4.0	3.2	8.3	10.4	17.6	13.8	27.1	44.5	9.9	43.6	-	3.5 3.5 0.43 0.70	6.1 C18	
Rectum	15 0	0.7	-	-	-	-	-	1.3	1.6	2.1	2.6	3.5	-	-	-	14.8	29.6	14.5	26.0	1.4 1.5 0.06 0.28	2.2 C19-20	
Anus	3 0	-	-	-	-	-	-	-	-	-	-	3.5	-	-	-	-	-	-	0.3 0.3 0.03 0.03	0.3 C21		
Liver	32 0	-	-	-	-	-	-	1.8	-	6.4	4.1	7.8	14.1	18.5	5.4	29.7	39.4	14.5	26.0	3.1 3.1 0.29 0.64	5.0 C22	
Gallbladder etc.	5 0	-	-	-	-	-	-	-	-	-	-	-	5.4	14.8	-	-	17.3	-	0.5 0.5 0.03 0.10	0.8 C23-24		
Pancreas	3 0	-	-	-	-	-	-	-	-	-	-	9.2	-	-	-	-	14.5	-	-	0.3 0.3 0.05 0.05	0.5 C25	
Nose, sinuses etc.	5 0	0.7	-	-	-	-	-	-	-	-	-	4.6	5.4	7.4	-	14.5	-	-	0.5 0.5 0.05 0.09	0.8 C30-31		
Larynx	26 0	-	-	-	-	-	-	-	-	-	2.6	-	18.5	37.9	29.7	19.7	43.6	43.3	-	2.5 2.5 0.30 0.54	4.6 C32	
Trachea, bronchus and lung	30 0	-	-	-	-	-	-	-	1.6	4.1	7.8	7.0	46.2	37.9	29.7	9.9	-	-	-	2.9 2.9 0.52 0.72	5.6 C33-34	
Other thoracic organs	1 0	-	-	-	-	-	-	-	-	-	4.6	-	-	-	-	-	-	-	0.1 0.1 0.02 0.02	0.2 C37-38		
Bone	23 0	-	-	1.7	1.7	0.9	5.6	1.3	3.2	4.1	-	3.5	4.6	16.3	7.4	-	-	17.3	-	2.2 2.2 0.21 0.25	2.8 C40-41	
Melanoma of skin	21 0	-	-	-	-	-	-	1.3	-	4.1	7.8	14.1	-	5.4	14.8	29.6	29.1	26.0	-	2.0 2.0 0.16 0.39	3.3 C43	
Other skin	54 0	-	-	-	-	2.5	5.5	-	9.4	4.8	4.1	5.2	14.1	4.6	16.3	51.9	49.3	43.6	69.3	5.2 5.2 0.33 0.84	7.3 C44	
Mesothelioma	5 0	-	-	-	-	-	-	-	-	-	2.6	-	-	10.8	7.4	9.9	-	-	-	0.5 0.5 0.07 0.15	1.0 C45	
Kaposi's sarcoma	155 0	0.8	1.7	0.8	-	5.6	49.7	51.3	66.2	39.1	35.1	13.8	27.1	22.2	19.7	43.6	34.7	-	14.8 15.0 1.46 1.67	18.4 C46		
Connective and soft tissue	23 0	2.0	-	3.4	0.9	1.1	5.4	-	6.2	2.6	7.0	4.6	10.8	-	-	-	-	8.7	-	2.2 2.2 0.22 0.22	2.6 C47-C49	
Breast	7 0	-	-	-	-	-	-	-	-	2.6	3.5	-	5.4	7.4	-	14.5	-	-	17.3	0.7 0.7 0.06 0.09	1.1 C50	
Penis	16 0	-	-	-	-	-	-	-	4.8	2.1	2.6	10.5	-	5.4	22.2	-	-	-	14.5	1.6 1.6 0.13 0.24	2.4 C60	
Prostate	222 0	-	-	-	-	-	-	-	4.0	1.6	2.1	13.0	35.1	78.5	135.5	222.5	325.2	639.5	459.4	21.2 21.5 1.35 0.40	3.57 C61	
Testis	10 0	2.0	-	-	-	0.8	1.8	-	-	-	4.1	-	3.5	-	-	-	-	-	14.5	1.0 1.0 0.06 0.06	1.0 C62	
Other male genital organs	2 0	-	-	-	-	0.8	-	-	-	-	-	-	-	-	-	-	-	-	0.2 0.2 0.00 0.05	0.3 C63		
Kidney	14 0	0.7	0.8	-	-	-	-	-	-	2.1	5.2	-	9.2	5.4	37.1	-	14.5	-	-	1.3 1.4 0.12 0.30	2.4 C64	
Renal pelvis	0 0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.0 0.0 0.00 0.00	0.0 C65		
Ureter	0 0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.0 0.0 0.00 0.00	0.0 C66		
Bladder	22 0	-	0.8	0.8	-	-	-	1.3	1.6	-	-	-	-	13.8	27.1	37.1	19.7	-	26.0	2.1 2.1 0.23 0.51	3.7 C67	
Other urinary organs	0 0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.0 0.0 0.00 0.00	0.0 C68		
Eye	54 0	1.3	-	0.8	2.8	1.1	4.0	30.5	26.9	5.2	10.5	9.2	5.4	14.8	9.9	-	-	8.7	-	5.2 5.2 0.49 0.61	6.4 C69	
Brain, nervous system	14 0	0.7	0.8	0.8	-	-	5.4	-	-	2.6	-	4.6	-	14.8	9.9	14.5	-	8.7	-	1.3 1.4 0.07 0.20	1.8 C70-72	
Thyroid	4 0	-	-	-	-	-	-	-	-	2.1	-	-	-	-	-	-	-	-	0.4 0.4 0.06 0.11	0.8 C73		
Adrenal gland	0 0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.0 0.0 0.00 0.00	0.0 C74		
Other endocrine	1 0	-	-	-	-	-	-	-	1.3	-	-	-	-	-	-	-	-	-	0.1 0.1 0.01 0.01	0.1 C75		
Hodgkin disease	10 0	-	-	2.4	-	0.8	-	1.1	1.3	-	2.1	-	3.5	4.6	-	-	-	-	14.5	1.0 1.0 0.08 0.08	1.1 C81	
Non-Hodgkin lymphoma	61 0	2.6	4.0	-	-	0.9	4.5	6.7	16.0	12.4	23.5	14.1	9.2	5.4	-	39.4	72.7	8.7	-	5.8 5.9 0.50 0.69	7.6 C82-85,C96	
Immunoproliferative diseases	0 0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.0 0.0 0.00 0.00	0.0 C88		
Multiple myeloma	9 0	-	-	-	-	-	-	-	-	1.6	-	2.6	7.0	-	5.4	14.8	9.9	14.5	-	0.9 0.9 0.08 0.21	1.6 C90	
Lymphoid leukaemia	13 0	0.7	3.2	0.8	0.8	0.9	-	-	-	-	2.6	-	5.4	-	9.9	14.5	-	-	8.7	1.2 1.3 0.07 0.12	1.7 C91-94	
Myeloid leukaemia	13 0	0.7	-	0.8	0.8	-	1.1	-	3.2	2.1	5.2	-	4.6	5.4	-	9.9	14.5	-	-	0.2 0.2 0.01 0.01	0.2 C95	
Leukaemia unspecified	2 0	-	-	0.8	-	0.8	-	-	-	-	-	-	-	-	-	-	-	-	0.2 0.2 0.01 0.01	0.2 C96		
Myeloproliferative disorders	3 0	-	-	-	-	-	-	-	1.3	-	2.6	-	-	-	-	-	-	-	0.3 0.3 0.02 0.06	0.5 MPD		
Myelodysplastic syndromes	0 0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.1 0.1 0.00 0.00	0.1 MDS		
Other and unspecified	68 0	-	0.8	0.8	0.8	1.8	3.2	7.7	6.1	9.5	4.5	22.6	14.7	30.9	16.4	21.5	39.6	64.5	-	6.1 5.2 0.52 0.71	7.2 O&U	
All sites	1362 2	12.1	8.2	11.4	6.6	25.1	38.6	149.8	195.2	224.1	270.1	390.2	401.2	636.2	518.4	780.7	633.6	488.5	-	122.8 11.8 18.63 16.30	ALL	
All sites but C44	1300 1	12.1	7.4	11.4	6.6	23.3	37.5	144.6	189.1	216.5	261.0	376.1	382.8	605.3	491.2	759.2	594.0	433.2	-	117.3 100.0 11.38 17.63	151.3 ALLbC44	

, Reference population: World Standard Population

Table builtFri Nov 18 07:56:36 SAST 2016 by CanReg5.

Central Cancer Registry, NAMIBIA (2013)

2013 Population Namibia

Incidence per 100,000 by age group (Period) - Female

SITE	ALL AGES UNK</
------	----------------

Central Cancer Registry, NAMIBIA (2014)

2014 Population Namibia

Incidence per 100,000 by age group (Period) - Male

S I T E	ALL AGES UNK	Incidence per 100,000 by age group (Period) - Male													CRUDE RATE	(%)	CUM 0-64	CUM 0-74	ASR	ICD (10th)							
		0-	5-	10-	15-	20-	25-	30-	35-	40-	45-	50-	55-	60-	65-	70-	75-	80+									
Lip	3 0	-	-	-	-	-	-	-	-	1.6	-	-	-	-	-	-	-	-	0.3	0.3 0.01 0.06	0.4 C00						
Tongue	31 0	-	-	-	-	-	-	-	-	-	2.6	31.6	27.7	27.1	7.4	39.4	14.5	34.7	3.0	2.9 0.45 0.68	5.4 C01-02						
Mouth	24 0	-	-	-	-	-	-	-	-	-	15.6	10.5	4.6	27.1	14.8	29.6	-	17.3	2.3	2.3 0.30 0.52	4.1 C03-06						
Salivary glands	11 0	-	0.8	0.8	-	0.9	-	-	-	1.6	8.3	-	-	4.6	-	-	-	17.3	1.1	1.0 0.09 0.09	1.2 C07-08						
Tonsil	5 0	-	-	-	0.8	-	-	-	-	-	-	3.5	4.6	5.4	-	-	-	-	0.5	0.5 0.07 0.12	0.9 C09						
Other oropharynx	3 0	-	-	-	-	-	-	-	-	-	-	3.5	-	-	14.8	-	-	-	0.3	0.3 0.02 0.09	0.6 C10						
Nasopharynx	6 0	-	-	-	0.8	-	0.9	-	-	1.6	-	-	4.6	5.4	-	-	-	8.7	0.6	0.6 0.07 0.07	0.7 C11						
Hypopharynx	2 0	-	-	-	-	-	-	-	-	-	-	9.2	-	-	-	-	-	-	0.2	0.2 0.05 0.05	0.4 C12-13						
Pharynx unspecified	2 0	-	-	-	-	-	-	-	-	-	-	3.5	-	-	9.9	-	-	-	0.2	0.2 0.02 0.07	0.4 C14						
Oesophagus	31 0	-	-	-	-	-	-	-	-	-	5.2	17.6	9.2	32.5	51.9	9.9	-	69.3	3.0	2.9 0.32 0.63	5.3 C15						
Stomach	16 0	-	-	-	-	-	-	-	-	2.2	1.3	1.6	-	2.6	7.0	13.8	5.4	7.4	9.9	14.5	1.5	1.5 0.17 0.26	2.4 C16				
Small intestine	1 0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.1	0.1 0.00 0.04	0.2 C17						
Colon	27 0	-	-	-	-	-	-	-	-	1.1	1.3	1.6	-	10.4	17.6	18.5	10.8	14.8	9.9	58.1	2.6	2.5 0.31 0.43	4.3 C18				
Rectum	15 0	-	-	-	-	-	-	-	-	0.9	2.2	1.3	1.6	-	7.8	7.0	4.6	10.8	-	-	14.5	1.4	1.4 0.18 0.22	2.2 C19-20			
Anus	6 0	-	-	-	-	-	-	-	-	-	-	2.1	2.6	-	-	-	-	-	0.6	0.6 0.02 0.12	0.8 C21						
Liver	17 0	-	-	-	-	-	-	-	-	1.1	-	-	2.1	2.6	-	-	-	-	1.6	1.6 0.15 0.34	2.8 C22						
Gallbladder etc.	2 0	-	-	-	-	-	-	-	-	-	-	-	-	2.6	-	-	-	-	0.2	0.2 0.04 0.04	0.4 C23-24						
Pancreas	5 0	-	-	-	-	-	-	-	-	-	-	-	-	3.5	4.6	5.4	14.8	-	-	-	0.5	0.5 0.07 0.14	1.0 C25				
Nose, sinuses etc.	5 0	-	-	-	-	-	-	-	-	1.8	1.1	-	-	-	3.5	-	5.4	-	-	-	-	0.5	0.5 0.06 0.06	0.6 C30-31			
Larynx	26 0	-	-	-	-	-	-	-	-	-	-	-	-	5.2	10.5	18.5	21.7	7.4	78.8	43.6	8.7	2.5	2.4 0.28 0.71	4.8 C32			
Trachea, bronchus and lung	42 0	-	-	-	-	-	-	-	-	-	-	-	-	2.1	5.2	3.5	41.5	10.8	81.6	88.7	29.1	4.0	3.9 0.32 1.17	7.7 C33-34			
Other thoracic organs	3 0	-	-	-	-	-	-	-	-	-	-	-	-	1.6	2.6	-	-	-	-	-	-	0.3	0.3 0.00 0.05	0.5 C37-38			
Bone	19 0	-	-	-	-	-	-	-	-	0.8	2.5	2.8	1.1	2.1	-	7.0	9.2	-	7.4	9.9	-	26.0	1.8 1.8 0.13	0.2 C24-41			
Melanoma of skin	16 0	-	-	-	-	-	-	-	-	-	1.3	3.2	4.1	2.6	3.5	4.6	5.4	7.4	9.9	29.1	26.0	1.5 1.5 0.12	0.2 C43				
Other skin	95 0	-	-	-	-	-	-	-	-	2.5	-	2.2	12.1	8.0	8.3	18.2	28.1	9.2	27.1	59.3	128.1	159.9	121.3	9.1 8.9 0.60	1.53 C44		
Mesothelioma	0 0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.0	0.0 0.00 0.00	0.0 C45			
Kaposi's sarcoma	160 0	-	0.7	-	-	-	-	-	-	4.2	0.9	11.2	30.9	43.3	35.1	52.1	56.2	32.3	48.8	22.2	78.8	58.1	78.0	15.3 15.0	1.58 2.08	2.8 C46	
Connective and soft tissue	23 0	-	0.7	2.4	-	-	-	-	-	1.7	0.9	2.2	8.1	-	2.1	7.0	-	-	9.9	43.6	8.7	-	-	2.2 2.2	0.13 0.17	2.4 C47,C49	
Breast	13 0	-	-	-	-	-	-	-	-	-	1.3	-	-	2.1	5.2	-	9.2	-	14.8	19.7	-	-	26.0	1.2	1.2 0.09	0.26 C50	
Penis	10 0	-	-	-	-	-	-	-	-	-	-	-	-	4.1	7.8	7.0	-	-	-	-	43.6	-	-	1.0	0.9 0.09	0.09 C51	
Prostate	266 1	-	-	-	-	-	-	-	-	-	-	-	-	2.6	31.6	115.4	103.0	348.6	443.4	755.8	580.7	-	25.4	25.0	1.27	5.24	43.3 C61
Testis	7 0	-	-	-	-	-	-	-	-	0.8	-	2.2	-	-	3.5	4.6	-	7.4	-	-	8.7	0.7	0.7 0.06	0.09 C62			
Other male genital organs	4 0	-	0.7	-	-	-	-	-	-	-	-	-	-	4.1	-	-	4.6	-	-	-	-	0.4	0.4 0.05	0.05 C63			
Kidney	26 0	-	4.6	2.4	-	-	-	-	-	2.5	-	1.3	-	10.3	2.6	3.5	9.2	-	14.8	9.9	-	-	-	2.5	2.4 0.18	0.31 C64	
Renal pelvis	0 0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.0	0.0 0.00	0.00 C65		
Ureter	0 0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.0	0.0 0.00	0.00 C66		
Bladder	20 0	-	-	-	-	-	-	-	-	-	-	-	-	3.2	2.1	5.2	14.1	9.2	10.8	22.2	29.6	14.5	-	-	1.9	1.9 0.22	0.48 C67
Other urinary organs	1 0	-	-	-	-	-	-	-	-	-	-	-	-	-	3.5	-	-	-	-	-	-	-	0.1	0.1 0.02	0.2 C68		
Eye	32 0	-	2.0	-	-	-	-	-	-	1.1	2.7	8.0	6.2	7.8	-	23.1	16.3	14.8	-	19.7	29.1	8.7	3.1	3.0 0.34	0.51 C69		
Brain, nervous system	14 0	-	-	1.6	1.7	2.5	-	-	1.1	1.3	1.6	4.1	2.6	-	4.1	-	-	-	-	-	-	-	1.3	1.3 0.11	0.11 C70-72		
Thyroid	4 0	-	-	-	-	-	-	-	-	-	2.2	1.3	-	3.8	-	-	-	-	-	-	-	0.4	0.4 0.04	0.04 C73			
Adrenal gland	0 0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.0	0.0 0.00	0.00 C74		
Other endocrine	1 0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.1	0.1 0.03	0.03 C75		
Hodgkin disease	6 0	-	-	0.8	0.8	-	-	-	-	1.1	-	-	-	2.1	-	-	-	-	-	-	-	0.6	0.6 0.05	0.05 C81			
Non-Hodgkin lymphoma	69 0	-	2.6	4.0	0.8	3.4	0.9	6.7	9.4	9.6	8.3	13.0	17.6	9.2	32.5	37.1	39.4	14.5	-	26.0	-	6.6 6.5	0.59 0.97	9.0 C82-85,C96			
Immunoproliferative diseases	2 0	-	-	-	-	-	-	-	-	-	-	-	-	2.1	-	-	-	-	-	-	-	0.2	0.2 0.01	0.01 C88			
Multiple myeloma	3 0	-	-	-	-	-	-	-	-	-	-	-	-	-	3.5	4.6	-	-	-	-	-	0.3	0.3 0.04	0.5 C90			
Lymphoid leukaemia	10 0	-	-	0.8	-	-	-	-	-	1.7	-	3.4	-	6.2	2.6	3.5	4.6	16.3	7.4	-	-	8.7	1.0	0.9 0.06	0.11 C91		
Myeloid leukaemia	18 0	-	0.7	0.8	-	-	-	-	-	1.7	-	3.4	-	6.2	2.6	3.5	4.6	16.3	7.4	-	-	8.7	1.7	1.7 0.20	0.24 C92-94		
Leukaemia unspecified	1 0	-	-	-	-	-	-	-	-	1.1	-	-	-	-	-	-	-	-	-	-	-	0.1	0.1 0.01	0.01 C95			
Myeloproliferative disorders	1 0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.1	0.1 0.03	0.03 MPD			
Myelodysplastic syndromes	2 0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.2	0.2 0.00	0.04 MDS			
Other and unspecified	53 0	-	0.7	-	0.8	0.8	-	-	-	2.7	3.2	6.2	7.8	36.9	21.7	51.9	49.3	72.7	34.7	-	-	5.1	5.0	0.53	1.03	8.6 O&U	
All sites	1159 1	12.6	12.9	9.1	22.8	15.7	45.9	77.9	93.1	124.0	195.4	358.4	470.8	477.0	882.5	1182.4	1482.6	1239.5	-	-	-	110.8	9.59	19.92	17.3 ALL		
All sites but C44	1064 1	12.6	12.9	6.6	22.8	12.0	43.7	65.8	85.0	115.8	177.2	330.3	461.6	449.9	823.2	1054.3	1322.7	1118.1	-	-	-	101.7	100.0	8.99	18.39	159.8 ALL	
																										ALL	

Central Cancer Registry, NAMIBIA (2010-2014)

CI5 Editorial Table 1
Number of cases in major diagnosis groups in single calendar years of observation

MALES

SITE	2010	2011	2012	2013	2014
Lip, oral cavity and pharynx (C00-14)	70 (7.1)	50 (5.8)	62 (7.3)	66 (6.4)	87 (8.2)
Digestive organs (C15-26)	128 (13)	100 (11.7)	94 (11.1)	132 (12.8)	121 (11.4)
Respiratory organs (C30-39)	91 (9.2)	50 (5.8)	57 (6.7)	62 (6)	76 (7.1)
Bone, cartilage, melanoma (C40-43)	30 (3)	28 (3.3)	23 (2.7)	44 (4.3)	35 (3.3)
Male genital (C60-63)	247 (25.1)	231 (27)	205 (24.1)	250 (24.2)	286 (26.9)
Urinary organs (C64-68)	24 (2.4)	25 (2.9)	31 (3.6)	36 (3.5)	44 (4.1)
Eye, brain, thyroid etc. (C69-75)	91 (9.2)	64 (7.5)	60 (7.1)	73 (7.1)	51 (4.8)
Haematopoietic (C81-96)	59 (6)	49 (5.7)	71 (8.4)	108 (10.5)	109 (10.3)
Other and unspecified (O&U)	227 (23.1)	242 (28.3)	238 (28)	253 (24.5)	241 (22.7)
All sites but C44 (ALLbC44)	984 (100)	855 (100)	850 (100)	1031 (100)	1063 (100)


FEMALES

SITE	2010	2011	2012	2013	2014
Lip, oral cavity and pharynx (C00-14)	50 (4.5)	30 (3.2)	34 (3.5)	44 (3.4)	50 (3.5)
Digestive organs (C15-26)	99 (8.9)	68 (7.2)	68 (7.1)	111 (8.5)	105 (7.2)
Respiratory organs (C30-39)	42 (3.8)	24 (2.5)	27 (2.8)	32 (2.5)	32 (2.2)
Bone, cartilage, melanoma (C40-43)	39 (3.5)	38 (4)	32 (3.3)	39 (3)	48 (3.3)
Breast (C50)	292 (26.2)	265 (28)	274 (28.5)	302 (23.2)	445 (30.7)
Female genital (C51-58)	296 (26.6)	230 (24.3)	230 (24)	389 (29.9)	410 (28.3)
Urinary organs (C64-68)	24 (2.2)	18 (1.9)	27 (2.8)	35 (2.7)	34 (2.3)
Eye, brain, thyroid etc. (C69-75)	64 (5.7)	71 (7.5)	85 (8.9)	77 (5.9)	75 (5.2)
Haematopoietic (C81-96)	43 (3.9)	50 (5.3)	54 (5.6)	81 (6.2)	85 (5.9)
Other and unspecified (O&U)	165 (14.8)	151 (16)	129 (13.4)	189 (14.5)	165 (11.4)
All sites but C44 (ALLbC44)	1114 (100)	945 (100)	960 (100)	1299 (100)	1449 (100)

BOTH SEXES

SITE	2010	2011	2012	2013	2014
Lip, oral cavity and pharynx (C00-14)	120 (5.7)	80 (4.4)	96 (5.3)	110 (4.7)	137 (5.4)
Digestive organs (C15-26)	227 (10.8)	168 (9.3)	162 (9)	243 (10.4)	226 (9)
Respiratory organs (C30-39)	133 (6.3)	74 (4.1)	84 (4.6)	94 (4)	108 (4.3)
Bone, cartilage, melanoma (C40-43)	69 (3.3)	66 (3.7)	55 (3)	83 (3.6)	83 (3.3)
Breast (C50)	309 (14.7)	281 (15.6)	283 (15.6)	309 (13.3)	458 (18.2)
Female genital (C51-58)	296 (14.1)	230 (12.8)	230 (12.7)	389 (16.7)	412 (16.4)
Male genital (C60-63)	247 (11.8)	231 (12.8)	205 (11.3)	250 (10.7)	286 (11.4)
Urinary organs (C64-68)	48 (2.3)	43 (2.4)	58 (3.2)	71 (3)	78 (3.1)
Eye, brain, thyroid etc. (C69-75)	155 (7.4)	135 (7.5)	145 (8)	150 (6.4)	126 (5)
Haematopoietic (C81-96)	102 (4.9)	99 (5.5)	125 (6.9)	189 (8.1)	195 (7.7)
Other and unspecified (O&U)	392 (18.7)	393 (21.8)	367 (20.3)	442 (19)	410 (16.3)
All sites but C44 (ALLbC44)	2098 (100)	1800 (100)	1810 (100)	2330 (100)	2519 (100)

**ALL SITES BUT SKIN
(BOTH SEXES)**



Central Cancer Registry, NAMIBIA (2010-2014)

CI5 Editorial Table 2
ASR in major diagnosis groups in single calendar years of observation

MALES

SITE	2010	2011	2012	2013	2014
Lip, oral cavity and pharynx (C00-14)	11.9 (7.8)	8 (6.3)	10.25 (7.9)	10.87 (7.2)	14.05 (8.8)
	22.18	15.95	15.79	21.33	19.72

Digestive organs (C15-26)	(14.4)	(12.6)	(12.2)	(14.1)	(12.3)
Respiratory organs (C30-39)	16.4 (10.7)	8.57 (6.8)	10.37 (8)	11.22 (7.4)	13.52 (8.5)
Bone, cartilage, melanoma (C40-43)	3.83 (2.5)	4.12 (3.3)	3.05 (2.4)	6.05 (4)	4.44 (2.8)
Male genital (C60-63)	41.25 (26.9)	38.22 (30.2)	35.03 (27.1)	39.42 (26.1)	46.1 (28.9)
Urinary organs (C64-68)	3.78 (2.5)	3.96 (3.1)	4.56 (3.5)	6.18 (4.1)	6.34 (4)
Eye, brain, thyroid etc. (C69-75)	11.99 (7.8)	8.02 (6.3)	7.81 (6.1)	8.99 (5.9)	6.68 (4.2)
Haematopoietic (C81-96)	8.86 (5.8)	6.73 (5.3)	9.51 (7.4)	13.56 (9)	14.18 (8.9)
Other and unspecified (O&U)	30.67 (20)	30.94 (24.4)	31.29 (24.2)	32.41 (21.4)	32.68 (20.5)
All sites but C44 (ALLbC44)	153.52 (100)	126.75 (100)	129.07 (100)	151.11 (100)	159.69 (100)

FEMALES


SITE	2010	2011	2012	2013	2014
Lip, oral cavity and pharynx (C00-14)	6.18 (4.6)	3.87 (3.5)	4.7 (4)	5.68 (3.6)	6.43 (3.7)
Digestive organs (C15-26)	12.88 (9.6)	8.46 (7.6)	8.43 (7.2)	13.58 (8.7)	13.75 (7.8)
Respiratory organs (C30-39)	6.05 (4.5)	3.12 (2.8)	3.58 (3.1)	4.47 (2.9)	4.48 (2.5)
Bone, cartilage, melanoma (C40-43)	4.21 (3.1)	4.04 (3.6)	3.38 (2.9)	4.6 (3)	5.55 (3.2)
Breast (C50)	37.43 (28)	33.29 (30)	35.43 (30.4)	38.66 (24.8)	55.85 (31.7)
Female genital (C51-58)	35.6 (26.6)	27.79 (25.1)	29.37 (25.2)	47.28 (30.3)	51.06 (29)
Urinary organs (C64-68)	3.02 (2.3)	1.8 (1.6)	2.97 (2.5)	4.37 (2.8)	4.42 (2.5)
Eye, brain, thyroid etc. (C69-75)	6.5 (4.9)	7.12 (6.4)	8.98 (7.7)	8.43 (5.4)	8.46 (4.8)
Haematopoietic (C81-96)	4.85 (3.6)	5.54 (5)	5.93 (5.1)	9.23 (5.9)	8.56 (4.9)
Other and unspecified (O&U)	17.07 (12.8)	15.77 (14.2)	13.71 (11.8)	19.57 (12.6)	17.41 (9.9)
All sites but C44 (ALLbC44)	133.78 (100)	110.8 (100)	116.49 (100)	155.86 (100)	175.96 (100)

BOTH SEXES

SITE	2010	2011	2012	2013	2014
Lip, oral cavity and pharynx (C00-14)	8.67 (6.1)	5.73 (4.9)	7.06 (5.8)	7.85 (5.2)	9.79 (5.9)
Digestive organs (C15-26)	16.85 (11.9)	11.71 (10)	11.61 (9.6)	16.94 (11.1)	16.22 (9.7)
Respiratory organs (C30-39)	10.57 (7.5)	5.5 (4.7)	6.5 (5.4)	7.41 (4.9)	8.37 (5)
Bone, cartilage, melanoma (C40-43)	4 (2.8)	4.04 (3.5)	3.19 (2.6)	5.25 (3.4)	5.11 (3.1)
Breast (C50)	21.83 (15.5)	19.37 (16.6)	20.18 (16.7)	21.8 (14.3)	31.7 (19)

Female genital (C51-58)	19.62 (13.9)	15.33 (13.1)	16.26 (13.4)	26.01 (17.1)	28.18 (16.9)
Male genital (C60-63)	17.43 (12.4)	15.98 (13.7)	14.78 (12.2)	16.6 (10.9)	19.24 (11.5)
Urinary organs (C64-68)	3.3 (2.3)	2.73 (2.3)	3.66 (3)	5.13 (3.4)	5.31 (3.2)
Eye, brain, thyroid etc. (C69-75)	8.96 (6.4)	7.47 (6.4)	8.41 (7)	8.68 (5.7)	7.6 (4.6)
Haematopoietic (C81-96)	6.68 (4.7)	6.09 (5.2)	7.52 (6.2)	11.28 (7.4)	11.05 (6.6)
Other and unspecified (O&U)	23.15 (16.4)	22.66 (19.4)	21.76 (18)	25.48 (16.7)	24.31 (14.6)
All sites but C44 (ALLbC44)	141.06 (100)	116.61 (100)	120.94 (100)	152.41 (100)	166.88 (100)

**ALL SITES BUT SKIN
(BOTH SEXES)**



Central Cancer Registry, NAMIBIA (2010–2014)

Data Quality Indicators

MALE

SITE	Cases	% Total	ASR(se)	MV(%)	CLIN(%)	DCO(%)	ICD10
Mouth & pharynx	335	6.54	11.01 (0.63)	99.40	0.60	0	C00–14
Oesophagus	108	2.11	3.87 (0.38)	92.59	7.41	0	C15
Stomach	72	1.41	2.32 (0.28)	100.00	0.00	0	C16
Colon, rectum, anus	221	4.32	7.05 (0.49)	98.64	1.36	0	C18–21
Liver	124	2.42	3.99 (0.37)	73.39	26.61	0	C22
Pancreas	28	0.55	1.04 (0.20)	60.71	39.29	0	C25
Larynx	147	2.87	5.30 (0.45)	100.00	0.00	0	C32
Lung, trachea, bronchus	161	3.14	5.88 (0.47)	78.88	21.12	0	C33–34
Pleura & other thoracic	8	0.16	0.22 (0.08)	100.00	0.00	0	C37–38
Melanoma of skin	88	1.72	2.62 (0.29)	100.00	0.00	0	C43
Prostate	1125	21.97	37.53 (1.17)	95.38	4.62	0	C61
Testis	32	0.62	0.70 (0.13)	96.88	3.12	0	C62
Kidney & urinary NOS	91	1.78	2.57 (0.29)	95.60	4.40	0	C64–66,68
Bladder	72	1.41	2.48 (0.31)	100.00	0.00	0	C67
Brain & nervous system	69	1.35	1.67 (0.22)	66.67	33.33	0	C70–72
Thyroid	21	0.41	0.65 (0.15)	100.00	0.00	0	C73
III-defined	188	3.67	6.09 (0.47)	98.40	1.60	0	C76–80
Lymphoma	289	5.64	7.85 (0.49)	97.92	2.08	0	C81–85,90,88,96
Leukaemia	107	2.09	2.71 (0.28)	100.00	0.00	0	C91–95
All sites but C44	4783	93.40	144.03 (2.20)	93.16	6.84	0	ALLbC44

FEMALE

SITE	Cases	% Total	ASR(se)	MV(%)	CLIN(%)	DCO(%)	ICD10
Mouth & pharynx	208	3.40	5.37 (0.39)	98.08	1.92	0	C00–14
Oesophagus	49	0.80	1.26 (0.19)	95.92	4.08	0	C15
Stomach	61	1.00	1.53 (0.21)	100.00	0.00	0	C16
Colon, rectum, anus	197	3.22	4.92 (0.36)	98.98	1.02	0	C18–21
Liver	81	1.32	2.04 (0.24)	86.42	13.58	0	C22
Pancreas	26	0.43	0.72 (0.15)	65.38	34.62	0	C25
Larynx	37	0.60	1.01 (0.17)	100.00	0.00	0	C32
Lung, trachea, bronchus	105	1.72	2.95 (0.30)	83.81	16.19	0	C33–34
Pleura & other thoracic	2	0.03	0.06 (0.04)	100.00	0.00	0	C37–38
Melanoma of skin	131	2.14	3.01 (0.28)	100.00	0.00	0	C43
Breast	1578	25.80	40.13 (1.05)	99.49	0.51	0	C50
Cervix	1116	18.24	27.11 (0.85)	98.84	1.16	0	C53
Corpus & Uterus NOS	153	2.50	4.12 (0.35)	99.35	0.65	0	C54–55
Ovary & adnexa	175	2.86	4.35 (0.34)	96.00	4.00	0	C56
Kidney & urinary NOS	80	1.31	1.99 (0.23)	97.50	2.50	0	C64–66,68
Bladder	60	0.98	1.40 (0.19)	100.00	0.00	0	C67
Brain & nervous system	77	1.26	1.70 (0.20)	77.92	22.08	0	C70–72
Thyroid	54	0.88	1.23 (0.17)	100.00	0.00	0	C73
III-defined	165	2.70	3.63 (0.30)	99.39	0.61	0	C76–80
Lymphoma	230	3.76	4.95 (0.34)	96.52	3.48	0	C81–85,90,88,96
Leukaemia	83	1.36	1.87 (0.22)	100.00	0.00	0	C91–95
All sites but C44	5767	94.28	138.58 (1.91)	96.76	3.24	0	ALLbC44

Cases of unknown age (8 M / 11 F) were excluded from these analyses

NOTES:

NOTES: